

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau(43) International Publication Date  
25 March 2004 (25.03.2004)

PCT

(10) International Publication Number  
WO 2004/024698 A1(51) International Patent Classification<sup>7</sup>: C07D 233/76,  
401/12, A61K 31/454D Alderley, Alderley Park, Macclesfield, Cheshire SK10  
4TG (GB).(21) International Application Number:  
PCT/GB2003/003937(74) Agent: ASTRAZENECA; Global Intellectual Property,  
P.O. Box 272, Mereside, Alderley Park, Macclesfield,  
Cheshire, SK10 4GR (GB).(22) International Filing Date:  
9 September 2003 (09.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0221250.4 13 September 2002 (13.09.2002) GB(71) Applicant (for all designated States except MG, US): AS-  
TRAZENECA AB [SE/SE]; Sodertalje, S-151 85 (SE).(71) Applicant (for MG only): ASTRAZENECA UK LIM-  
ITED [GB/GB]; 15 Stanhope Gate, London, Greater Lon-  
don W1K 1LN (GB).

(72) Inventors; and


(75) Inventors/Applicants (for US only): BURROWS,  
Jeremy, Nicholas [GB/GB]; AstraZeneca R & D Alder-  
ley, Alderley Park, Macclesfield, Cheshire SK10 4TG  
(GB). TUCKER, Howard [GB/GB]; AstraZeneca R &(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: 1-SULFONYL PIPERIDINE DERIVATIVES



(1)

(57) Abstract: Piperidine derivatives of formula (1) that are useful in the inhibition of metalloproteinases, in particular TNF- $\alpha$  Converting Enzyme (TACE) and thus in the treatment of autoimmune disease, allergic/atopic diseases, transplant rejection, graft versus host disease, cardiovascular disease, reperfusion injury and malignancy.

WO 2004/024698 A1

BEST AVAILABLE COPY

BEST AVAILABLE COPY

**1-ALPHONYL PIPERIDINE DERIVATIVES AS  
TACE SELECTIVE INHIBITORS**

The present invention relates to compounds useful in the inhibition of metalloproteinases and in particular to pharmaceutical compositions comprising these, as well as their use.

- 5       The compounds of this invention are inhibitors of one or more metalloproteinase enzymes and are particularly effective as inhibitors of TNF- $\alpha$  (Tumour Necrosis Factor- $\alpha$ ) Production. Metalloproteinases are a superfamily of proteinases (enzymes) whose numbers in recent years have increased dramatically. Based on structural and functional considerations these enzymes have been classified into families and subfamilies as described in N.M. Hooper
- 10 (1994) FEBS Letters 354:1-6. Examples of metalloproteinases include the matrix metalloproteinases (MMP) such as the collagenases (MMP1, MMP8, MMP13), the gelatinases (MMP2, MMP9), the stromelysins (MMP3, MMP10, MMP11), matrilysin (MMP7), metalloelastase (MMP12), enamelysin (MMP19), the MT-MMPs (MMP14, MMP15, MMP16, MMP17); the reprolysin or adamalysin or MDC family which includes the
- 15 secretases and sheddases such as TNF- $\alpha$  converting enzymes (ADAM10 and TACE); the ADAM-TS family (for example ADAM-TS1 and ADAM-TS4); the astacin family which include enzymes such as procollagen processing proteinase (PCP); and other metalloproteinases such as the endothelin converting enzyme family and the angiotensin converting enzyme family.
- 20       Metalloproteinases are believed to be important in a plethora of physiological disease processes that involve tissue remodelling such as embryonic development, bone formation and uterine remodelling during menstruation. This is based on the ability of the metalloproteinases to cleave a broad range of matrix substrates such as collagen, proteoglycan and fibronectin. Metalloproteinases are also believed to be important in the processing, or secretion, of
- 25 biologically important cell mediators, such as tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ); and the post translational proteolysis processing, or shedding, of biologically important membrane proteins, such as the low affinity IgE receptor CD23 (for a more complete list see N. M. Hooper *et al.*, (1997) Biochem J. 321:265-279).

- 30       Metalloproteinases have been associated with many disease conditions. Inhibition of the activity of one or more metalloproteinases may well be of benefit in these disease conditions, for example: various inflammatory and allergic diseases such as, inflammation of the joint (especially rheumatoid arthritis, osteoarthritis and gout), inflammation of the gastro-

intestinal tract (especially inflammatory bowel disease, ulcerative colitis and gastritis), inflammation of the skin (especially psoriasis, eczema and dermatitis); in tumour metastasis or invasion; in disease associated with uncontrolled degradation of the extracellular matrix such as osteoarthritis; in bone resorptive disease (such as osteoporosis and Paget's disease); in diseases associated with aberrant angiogenesis; the enhanced collagen remodelling associated with diabetes, periodontal disease (such as gingivitis), corneal ulceration, ulceration of the skin, post-operative conditions (such as colonic anastomosis) and dermal wound healing; demyelinating diseases of the central and peripheral nervous systems (such as multiple sclerosis); Alzheimer's disease; and extracellular matrix remodelling observed in cardiovascular diseases such as restenosis and atherosclerosis.

A number of metalloproteinase inhibitors are known; different classes of compounds may have different degrees of potency and selectivity for inhibiting various metalloproteinases. We have discovered a class of compounds that are inhibitors of metalloproteinases and are of particular interest in inhibiting TACE. The compounds of this invention have beneficial potency and/or pharmacokinetic properties.

TACE (also known as ADAM17) which has been isolated and cloned [R.A. Black *et al.* (1997) *Nature* 385:729-733; M.L. Moss *et al.* (1997) *Nature* 385:733-736] is a member of the adamalysin family of metalloproteins. TACE has been shown to be responsible for the cleavage of pro-TNF- $\alpha$ , a 26kDa-membrane bound protein to release 17kDa biologically active soluble TNF- $\alpha$  [Schlondorff *et al.* (2000) *Biochem. J.* 347: 131-138]. TACE mRNA is found in most tissues, however TNF- $\alpha$  is produced primarily by activated monocytes, macrophages and T lymphocytes. TNF- $\alpha$  has been implicated in a wide range of pro-inflammatory biological processes including induction of adhesion molecules and chemokines to promote cell trafficking, induction of matrix destroying enzymes, activation of fibroblasts to produce prostaglandins and activation of the immune system [Aggarwal *et al.* (1996) *Eur. Cytokine Netw.* 7: 93-124]. Clinical use of the anti-TNF biologicals has shown TNF- $\alpha$  to play an important role in a range of inflammatory diseases including rheumatoid arthritis, Crohn's disease and psoriasis [Onrust *et al.* (1998) *Biodrugs* 10: 397-422, Jarvis *et al.* (1999) *Drugs* 57:945-964]. TACE activity has also been implicated in the shedding of other membrane bound proteins including TGF $\alpha$ , p75 & p55 TNF receptors, L-selectin and amyloid precursor protein [Black (2002) *Int. J. Biochem. Cell Biol.* 34: 1-5]. The biology of TACE inhibition has recently been reviewed and shows TACE to have a central role in TNF- $\alpha$


production and selective TACE inhibitors to have equal, and possibly greater, efficacy in the collagen induced arthritis model of RA than strategies that directly neutralise TNF- $\alpha$  [Newton et al (2001) Ann. Rheum. Dis. 60: iii25-iii32].

A TACE inhibitor might therefore be expected to show efficacy in all disease where TNF- $\alpha$  has been implicated including, but not limited to, inflammatory diseases including rheumatoid arthritis and psoriasis, autoimmune diseases, allergic/atopic diseases, transplant rejection and graft versus host disease, cardiovascular disease, reperfusion injury, malignancy and other proliferative diseases. A TACE inhibitor might also be effective against respiratory disease such as asthma and chronic obstructive pulmonary diseases (referred to herein as COPD).

Metalloproteinase inhibitors are known in the art. WO 02/074750 and WO 02/074767 disclose compounds comprising a metal binding group that are inhibitors of metalloproteinases. WO 02/074751 also disclosed compounds that are inhibitors of metalloproteinases and especially MMP12.

We are able to provide further compounds that have metalloproteinase inhibitory activity, and are in particular inhibitors of TACE (ADAM17).

According to the first aspect of the present invention there is provided a compound of formula (1), a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof:



formula (1)

wherein:

Y<sup>1</sup> and Y<sup>2</sup> are independently O or S;

z is NR<sup>8</sup>, O or S;

n is 0 or 1;

W is NR<sup>1</sup>, CR<sup>1</sup>R<sup>2</sup> or a bond;

m is 0 or 1;

D is hydrogen, C<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl or fluoro;



X is  $-(CR^{12}R^{13})_t-Q-(CR^{14}R^{15})_u-$  where t and u are independently 0 or 1 and Q is O, S, SO or SO<sub>2</sub>;

B is a group selected from aryl, heteroaryl and heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl,

- 5 trifluoromethoxy, halo, cyano, C<sub>1-4</sub>alkyl (optionally substituted by R<sup>9</sup> or C<sub>1-4</sub>alkoxy or one or more halo), C<sub>2-4</sub>alkenyl (optionally substituted by halo or R<sup>9</sup>), C<sub>2-4</sub>alkynyl (optionally substituted by halo or R<sup>9</sup>), C<sub>3-6</sub>cycloalkyl (optionally substituted by R<sup>9</sup> or one or more halo), C<sub>5-6</sub>cycloalkenyl (optionally substituted by halo or R<sup>9</sup>), aryl (optionally substituted by halo or C<sub>1-4</sub>alkyl), heteroaryl (optionally substituted by halo or C<sub>1-4</sub>alkyl), heterocyclyl (optionally substituted by C<sub>1-4</sub>alkyl), -SR<sup>11</sup>, -SOR<sup>11</sup>, -SO<sub>2</sub>R<sup>11</sup>, -SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, -NR<sup>9</sup>SO<sub>2</sub>R<sup>11</sup>, -NHCONR<sup>9</sup>R<sup>10</sup>, -OR<sup>9</sup>, -NR<sup>9</sup>R<sup>10</sup>, -CONR<sup>9</sup>R<sup>10</sup> and -NR<sup>9</sup>COR<sup>10</sup>; or B is C<sub>2-4</sub>alkenyl or C<sub>2-4</sub>alkynyl, each being optionally substituted by a group selected from C<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl, aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more halo, nitro, cyano, trifluoromethyl, trifluoromethoxy, -CONHR<sup>9</sup>, -CONR<sup>9</sup>R<sup>10</sup>, -SO<sub>2</sub>R<sup>11</sup>, -SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, -NR<sup>9</sup>SO<sub>2</sub>R<sup>11</sup>, C<sub>1-4</sub>alkyl or C<sub>1-4</sub>alkoxy; with the provisos that:

- when n is 1 and W is NR<sup>1</sup>, CR<sup>1</sup>R<sup>2</sup> or a bond; or when n is 0 and W is CR<sup>1</sup>R<sup>2</sup>; then B is a group selected from aryl, heteroaryl and heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo, cyano, C<sub>1-4</sub>alkyl (optionally substituted by R<sup>9</sup> or C<sub>1-4</sub>alkoxy or one or more halo), C<sub>2-4</sub>alkenyl (optionally substituted by halo or R<sup>9</sup>), C<sub>2-4</sub>alkynyl (optionally substituted by halo or R<sup>9</sup>), C<sub>3-6</sub>cycloalkyl (optionally substituted by R<sup>9</sup> or one or more halo), C<sub>5-6</sub>cycloalkenyl (optionally substituted by halo or R<sup>9</sup>), aryl (optionally substituted by halo or C<sub>1-4</sub>alkyl), heteroaryl (optionally substituted by halo or C<sub>1-4</sub>alkyl), heterocyclyl (optionally substituted by C<sub>1-4</sub>alkyl), -SR<sup>11</sup>, -SOR<sup>11</sup>, -SO<sub>2</sub>R<sup>11</sup>, -SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, -NR<sup>9</sup>SO<sub>2</sub>R<sup>11</sup>, -NHCONR<sup>9</sup>R<sup>10</sup>, -OR<sup>9</sup>, -NR<sup>9</sup>R<sup>10</sup>, -CONR<sup>9</sup>R<sup>10</sup> and -NR<sup>9</sup>COR<sup>10</sup>; or B is C<sub>2-4</sub>alkenyl or C<sub>2-4</sub>alkynyl, each being optionally substituted by a group selected from C<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl, aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more halo, nitro, cyano, trifluoromethyl, trifluoromethoxy, -CONHR<sup>9</sup>, -CONR<sup>9</sup>R<sup>10</sup>, -SO<sub>2</sub>R<sup>11</sup>, -SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, -NR<sup>9</sup>SO<sub>2</sub>R<sup>11</sup>, C<sub>1-4</sub>alkyl or C<sub>1-4</sub>alkoxy; and
- 30 when n is 0 and W is NR<sup>1</sup> or a bond; then B is a group selected from bicyclic aryl, bicyclic heteroaryl and bicyclic heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo,

- cyano, C<sub>1-4</sub>alkyl (optionally substituted by R<sup>9</sup> or C<sub>1-4</sub>alkoxy or one or more halo), C<sub>2-4</sub>alkenyl (optionally substituted by halo or R<sup>9</sup>), C<sub>2-4</sub>alkynyl (optionally substituted by halo or R<sup>9</sup>), C<sub>3-6</sub>cycloalkyl (optionally substituted by R<sup>9</sup> or one or more halo), C<sub>5-6</sub>cycloalkenyl (optionally substituted by halo or R<sup>9</sup>), aryl (optionally substituted by halo or C<sub>1-4</sub>alkyl), heteroaryl
- 5 (optionally substituted by halo or C<sub>1-4</sub>alkyl), heterocyclyl (optionally substituted by C<sub>1-4</sub>alkyl), -SR<sup>11</sup>, -SOR<sup>11</sup>, -SO<sub>2</sub>R<sup>11</sup>, -SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, -NR<sup>9</sup>SO<sub>2</sub>R<sup>11</sup>, -NHCONR<sup>9</sup>R<sup>10</sup>, -OR<sup>9</sup>, -NR<sup>9</sup>R<sup>10</sup>, -CONR<sup>9</sup>R<sup>10</sup> and -NR<sup>9</sup>COR<sup>10</sup>; or B is C<sub>2-4</sub>alkenyl or C<sub>2-4</sub>alkynyl, each being optionally substituted by a group selected from C<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl, aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more halo, nitro, cyano, trifluoromethyl,
- 10 trifluoromethoxy, -CONHR<sup>9</sup>, -CONR<sup>9</sup>R<sup>10</sup>, -SO<sub>2</sub>R<sup>11</sup>, -SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, -NR<sup>9</sup>SO<sub>2</sub>R<sup>11</sup>, C<sub>1-4</sub>alkyl or C<sub>1-4</sub>alkoxy;
- R<sup>1</sup> and R<sup>2</sup> are independently hydrogen or a group selected from C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-6</sub>cycloalkyl and C<sub>5-6</sub>cycloalkenyl which group may be optionally substituted by halo, cyano, hydroxy or C<sub>1-4</sub>alkoxy;
- 15 R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are independently hydrogen or a group selected from C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-6</sub>cycloalkyl, C<sub>5-6</sub>cycloalkenyl, aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more substituents independently selected from halo, nitro, cyano, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>3-6</sub>cycloalkyl (optionally substituted by one or more R<sup>17</sup>), aryl (optionally substituted by one or more R<sup>17</sup>),
- 20 heteroaryl (optionally substituted by one or more R<sup>17</sup>), heterocyclyl, -OR<sup>18</sup>, -SR<sup>19</sup>, -SOR<sup>19</sup>, -SO<sub>2</sub>R<sup>19</sup>, -COR<sup>19</sup>, -CO<sub>2</sub>R<sup>18</sup>, -CONR<sup>18</sup>R<sup>20</sup>, -NR<sup>16</sup>COR<sup>18</sup>, -SO<sub>2</sub>NR<sup>18</sup>R<sup>20</sup> and -NR<sup>16</sup>SO<sub>2</sub>R<sup>19</sup>; or R<sup>1</sup> and R<sup>3</sup> together with the nitrogen or carbon atoms and carbon atom to which they are respectively attached form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and SO<sub>2</sub> where the ring is optionally
- 25 substituted on carbon by C<sub>1-4</sub>alkyl, fluoro or C<sub>1-4</sub>alkoxy and/or on nitrogen by -COC<sub>1-3</sub>alkyl, -SO<sub>2</sub>C<sub>1-3</sub>alkyl or C<sub>1-4</sub>alkyl;
- or R<sup>3</sup> and R<sup>4</sup> together form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and SO<sub>2</sub> where the ring is optionally substituted on carbon by C<sub>1-4</sub>alkyl, fluoro or C<sub>1-4</sub>alkoxy and/or on nitrogen by -COC<sub>1-3</sub>alkyl, -
- 30 SO<sub>2</sub>C<sub>1-3</sub>alkyl or C<sub>1-4</sub>alkyl;
- or R<sup>5</sup> and R<sup>6</sup> together form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and SO<sub>2</sub> where the ring is optionally

- substituted on carbon by  $C_{1-4}$ alkyl, fluoro or  $C_{1-4}$ alkoxy and/or on nitrogen by  $-COC_{1-3}$ alkyl,  $-SO_2C_{1-3}$ alkyl or  $C_{1-4}$ alkyl;
- $R^7$  is hydrogen or a group selected from  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl, heteroalkyl,  $C_{3-7}$ cycloalkyl, aryl, heteroaryl or heterocyclyl where the group is optionally substituted by halo,
- 5  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $C_{3-7}$ cycloalkyl, heterocyclyl, aryl, heteroaryl or heteroalkyl; and wherein the group from which  $R^7$  may be selected is optionally substituted on the group and/or on its optional substituent by one or more substituents independently selected from halo, cyano,  $C_{1-4}$ alkyl, nitro, halo $C_{1-4}$ alkyl, heteroalkyl, aryl, heteroaryl, hydroxy $C_{1-4}$ alkyl,  $C_{3-7}$ cycloalkyl, heterocyclyl,  $C_{1-4}$ alkoxy $C_{1-4}$ alkyl, halo $C_{1-4}$ alkoxy $C_{1-4}$ alkyl,  $-COC_{1-4}$ alkyl,  $-OR^{21}$ ,  $-CO_2R^{21}$ ,
- 10  $SR^{25}$ ,  $-SOR^{25}$ ,  $-SO_2R^{25}$ ,  $-NR^{21}COR^{22}$ ,  $-CONR^{21}R^{22}$  and  $-NHCONR^{21}R^{22}$ ;
- or  $R^3$  and  $R^7$  together with the carbon atoms to which they are each attached and  $(CR^5R^6)_n$  form a saturated 5- to 7-membered ring optionally containing a heteroatom group selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted on carbon by  $C_{1-4}$ alkyl, fluoro or  $C_{1-4}$ alkoxy and/or on nitrogen by  $-COC_{1-3}$ alkyl,  $-SO_2C_{1-3}$ alkyl or  $C_{1-4}$ alkyl;
- 15  $R^8$  is selected from hydrogen,  $C_{1-6}$ alkyl and halo $C_{1-6}$ alkyl;
- $R^9$  and  $R^{10}$  are independently hydrogen,  $C_{1-6}$ alkyl or  $C_{3-6}$ cycloalkyl;
- or  $R^9$  and  $R^{10}$  together with the nitrogen to which they are attached form a heterocyclic 4 to 7-membered ring.
- $R^{11}$  is  $C_{1-6}$ alkyl or  $C_{3-6}$ cycloalkyl;
- 20  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$  and  $R^{15}$  are independently selected from hydrogen,  $C_{1-6}$ alkyl and  $C_{3-6}$ cycloalkyl;
- $R^{16}$  is hydrogen or  $C_{1-6}$ alkyl;
- $R^{17}$  is selected from halo,  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl and  $C_{1-6}$ alkoxy;
- $R^{18}$  is hydrogen or a group selected from  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl,  $C_{3-7}$ cycloalkenyl, saturated heterocyclyl, aryl, heteroaryl, aryl $C_{1-4}$ alkyl and heteroaryl $C_{1-4}$ alkyl which group is optionally
- 25 substituted by one or more halo;
- $R^{19}$  and  $R^{25}$  are independently a group selected from  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl,  $C_{3-7}$ cycloalkenyl, saturated heterocyclyl, aryl, heteroaryl, aryl $C_{1-4}$ alkyl and heteroaryl $C_{1-4}$ alkyl which group is optionally substituted by one or more halo;
- $R^{20}$  is hydrogen,  $C_{1-6}$ alkyl or  $C_{3-6}$ cycloalkyl;
- 30 or  $R^{18}$  and  $R^{20}$  together with the nitrogen to which they are attached form a heterocyclic 4- to 7- membered ring;
- $R^{21}$  and  $R^{22}$  are independently hydrogen,  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl, aryl and aryl $C_{1-4}$ alkyl;

or  $R^{21}$  and  $R^{22}$  together with the nitrogen to which they are attached form a heterocyclic 5- to 6- membered ring.

According to a second aspect of the invention there is provided a compound of formula (1), a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof wherein:

- 5  $Y^1$  and  $Y^2$  are independently O or S;  
 $z$  is  $NR^8$ , O or S;  
 $n$  is 0;  
 $W$  is  $NR^1$  or a bond;  
 $m$  is 0 or 1;
- 10  $D$  is hydrogen,  $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl or fluoro;  
 $X$  is  $-(CR^{12}R^{13})_t-Q-(CR^{14}R^{15})_u-$  where  $t$  and  $u$  are independently 0 or 1 and  $Q$  is O, S, SO or  $SO_2$ ;  
 $B$  is a group selected from aryl, heteroaryl and heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl,
- 15 trifluoromethoxy, halo, cyano,  $C_{1-4}$ alkyl (optionally substituted by  $R^9$  or  $C_{1-4}$ alkoxy or one or more halo),  $C_{2-4}$ alkenyl (optionally substituted by halo or  $R^9$ ),  $C_{2-4}$ alkynyl (optionally substituted by halo or  $R^9$ ),  $C_{3-6}$ cycloalkyl (optionally substituted by  $R^9$  or one or more halo),  $C_{5-6}$ cycloalkenyl (optionally substituted by halo or  $R^9$ ), aryl (optionally substituted by halo or  $C_{1-4}$ alkyl), heteroaryl (optionally substituted by halo or  $C_{1-4}$ alkyl), heterocyclyl (optionally
- 20 substituted by  $C_{1-4}$ alkyl),  $-SR^{11}$ ,  $-SOR^{11}$ ,  $-SO_2R^{11}$ ,  $-SO_2NR^9R^{10}$ ,  $-NR^9SO_2R^{11}$ ,  $-NHCONR^9R^{10}$ ,  $-OR^9$ ,  $-CONR^9R^{10}$  and  $-NR^9COR^{10}$ ;  
 $R^1$  is hydrogen or a group selected from  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-6}$ cycloalkyl and  $C_{5-6}$ cycloalkenyl which group may be optionally substituted by halo, cyano, hydroxy or  $C_{1-4}$ alkoxy;
- 25  $R^3$  and  $R^4$  are independently hydrogen or a group selected from  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{3-5}$ cycloalkyl, pentenyl, aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more substituents independently selected from halo, nitro, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{3-6}$ cycloalkyl (optionally substituted by one or more  $R^{17}$ ), aryl (optionally substituted by one or more  $R^{17}$ ),
- 30 heteroaryl (optionally substituted by one or more  $R^{17}$ ), heterocyclyl,  $-OR^{18}$ ,  $-SR^{19}$ ,  $-SOR^{19}$ ,  $-SO_2R^{19}$ ,  $-CONR^{18}R^{20}$  and  $-NR^{16}COR^{18}$ ;

- or  $R^1$  and  $R^3$  together with the nitrogen and carbon atoms to which they are respectively attached form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted on carbon by  $C_{1-4}$ alkyl, fluoro or  $C_{1-4}$ alkoxy and/or on nitrogen by  $-COC_{1-3}$ alkyl,  $-SO_2C_{1-3}$ alkyl or  $C_{1-4}$ alkyl;
- or  $R^3$  and  $R^4$  together form a carbocyclic or saturated heterocyclic 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted on carbon by  $C_{1-4}$ alkyl, fluoro or  $C_{1-4}$ alkoxy and/or on nitrogen by  $-COC_{1-3}$ alkyl,  $-SO_2C_{1-3}$ alkyl or  $C_{1-4}$ alkyl;
- 10  $R^7$  is hydrogen or a group selected from  $C_{1-4}$ alkyl, heteroalkyl,  $C_{3-5}$ cycloalkyl, aryl, heteroaryl or heterocyclyl where the group is optionally substituted by halo,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $C_{3-5}$ cycloalkyl, heterocyclyl, aryl, heteroaryl or heteroalkyl; and wherein the group from which  $R^7$  may be selected is optionally substituted on the group and/or on its optional substituent by one or more substituents independently selected from halo, cyano,  $C_{1-4}$ alkyl, nitro, halo $C_{1-4}$ alkyl,
- 15 heteroalkyl, aryl, heteroaryl, hydroxy $C_{1-4}$ alkyl,  $C_{3-5}$ cycloalkyl, heterocyclyl,  $C_{1-4}$ alkoxy $C_{1-4}$ alkyl, halo $C_{1-4}$ alkoxy $C_{1-4}$ alkyl,  $-COC_{1-4}$ alkyl,  $-OR^{21}$ ,  $-CO_2R^{21}$ ,  $-SR^{25}$ ,  $-SOR^{25}$ ,  $-SO_2R^{25}$ ,  $-CONR^{21}R^{22}$  and  $-NHCONR^{21}R^{22}$ ;
- or  $R^3$  and  $R^7$  together with the carbon atoms to which they are each attached and  $(CR^5R^6)_n$  form a saturated carbocyclic or heterocyclic 5- or 6-membered ring;
- 20  $R^8$  is selected from hydrogen,  $C_{1-4}$ alkyl and halo $C_{1-4}$ alkyl;
- $R^9$  and  $R^{10}$  are independently hydrogen,  $C_{1-6}$ alkyl or  $C_{3-6}$ cycloalkyl;
- or  $R^9$  and  $R^{10}$  together with the nitrogen to which they are attached form a heterocyclic 4 to 6-membered ring;
- $R^{11}$  is  $C_{1-4}$ alkyl or  $C_{3-5}$ cycloalkyl;
- 25  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$  and  $R^{15}$  are independently selected from hydrogen,  $C_{1-4}$ alkyl and  $C_{3-4}$ cycloalkyl;
- $R^{16}$  is hydrogen or  $C_{1-4}$ alkyl;
- $R^{17}$  is selected from halo,  $C_{1-4}$ alkyl,  $C_{3-5}$ cycloalkyl and  $C_{1-4}$ alkoxy;
- $R^{18}$  is hydrogen or a group selected from  $C_{1-4}$ alkyl,  $C_{3-5}$ cycloalkyl,  $C_{5-6}$ cycloalkenyl, saturated heterocyclyl, aryl, heteroaryl, aryl $C_{1-4}$ alkyl and heteroaryl $C_{1-4}$ alkyl which group is optionally
- 30 substituted by one or more halo;

$R^{19}$  and  $R^{25}$  are independently a group selected from  $C_{1-4}$ alkyl,  $C_{3-5}$ cycloalkyl,  $C_5$ - $\epsilon$ cycloalkenyl, saturated heterocyclyl, aryl, heteroaryl, aryl $C_{1-4}$ alkyl and heteroaryl $C_{1-4}$ alkyl which group is optionally substituted by one or more halo;

$R^{20}$  is hydrogen,  $C_{1-4}$ alkyl or  $C_{3-5}$ cycloalkyl;

5 or  $R^{18}$  and  $R^{20}$  together with the nitrogen to which they are attached form a heterocyclic 4- to 6- membered ring;

$R^{21}$  and  $R^{22}$  are independently hydrogen,  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl, aryl and aryl $C_{1-4}$ alkyl;

or  $R^{21}$  and  $R^{22}$  together with the nitrogen to which they are attached form a heterocyclic 5- to 6- membered ring.

10 In another aspect of the present invention there is provided a compound of formula (1) or a pharmaceutically acceptable salt thereof.

It is to be understood that, insofar as certain of the compounds of formula (1) defined above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon or sulphur atoms, the invention includes in its definition any such optically active or  
15 racemic form which possesses metalloproteinases inhibition activity and in particular TACE inhibition activity. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, the above-mentioned activity may be evaluated using the standard laboratory techniques referred to hereinafter.

20 Compounds of formula (1) are therefore provided as enantiomers, diastereomers, geometric isomers and atropisomers.

Within the present invention it is to be understood that a compound of formula (1) or a salt thereof may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be  
25 understood that the invention encompasses any tautomeric form which has metalloproteinases inhibition activity and in particular TACE inhibition activity and is not to be limited merely to any one tautomeric form utilised within the formulae drawings.

It is also to be understood that certain compounds of formula (1) and salts thereof can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be  
30 understood that the invention encompasses all such solvated forms which have metalloproteinases inhibition activity and in particular TACE inhibition activity.



It is also to be understood that certain compounds of formula (1) may exhibit polymorphism, and that the invention encompasses all such forms which possess metalloproteinases inhibition activity and in particular TACE inhibition activity.

The present invention relates to compounds of formula (1) as defined herein as well  
5 as to the salts thereof. Salts for use in pharmaceutical compositions will be pharmaceutically acceptable salts, but other salts may be useful in the production of compounds of formula (1) and their pharmaceutically acceptable salts. Pharmaceutically acceptable salts of the invention may, for example, include acid addition salts of compounds of formula (1) as defined herein which are sufficiently basic to form such salts. Such acid addition salts include but are not  
10 limited to hydrochloride, hydrobromide, citrate and maleate salts and salts formed with phosphoric and sulphuric acid. In addition where compounds of formula (1) are sufficiently acidic, salts are base salts and examples include but are not limited to, an alkali metal salt for example sodium or potassium, an alkaline earth metal salt for example calcium or magnesium, or organic amine salts for example triethylamine or tris-(2-hydroxyethyl)amine

15 The compounds of formula (1) may also be provided as *in vivo* hydrolysable esters. An *in vivo* hydrolysable ester of a compound of formula (1) containing a carboxy or hydroxy group is, for example a pharmaceutically acceptable ester which is cleaved in the human or animal body to produce the parent acid or alcohol. Such esters can be identified by administering, for example, intravenously to a test animal, the compound under test and  
20 subsequently examining the test animal's body fluid.

Suitable pharmaceutically acceptable esters for carboxy include C<sub>1-6</sub>alkoxymethyl esters for example methoxymethyl, C<sub>1-6</sub>alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C<sub>3-8</sub>cycloalkoxycarbonyloxyC<sub>1-6</sub>alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example  
25 5-methyl-1,3-dioxolen-2-onylmethyl; and C<sub>1-6</sub>alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

Suitable pharmaceutically-acceptable esters for hydroxy include inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and  $\alpha$ -acyloxyalkyl ethers and  
30 related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of  $\alpha$ -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxymethoxy. A selection of *in vivo* hydrolysable ester forming groups



for hydroxy include C<sub>1-10</sub>alkanoyl, for example formyl, acetyl; benzoyl; phenylacetyl; substituted benzoyl and phenylacetyl, C<sub>1-10</sub>alkoxycarbonyl (to give alkyl carbonate esters), for example ethoxycarbonyl; di-(C<sub>1-4</sub>)alkylcarbamoyl and *N*-(di-(C<sub>1-4</sub>)alkylaminoethyl)-*N*-(C<sub>1-4</sub>)alkylcarbamoyl (to give carbamates); di-(C<sub>1-4</sub>)alkylaminoacetyl and carboxyacetyl.

- 5 Examples of ring substituents on phenylacetyl and benzoyl include aminomethyl, (C<sub>1-4</sub>)alkylaminomethyl and di-((C<sub>1-4</sub>)alkyl)aminomethyl, and morpholino or piperazino linked from a ring nitrogen atom via a methylene linking group to the 3- or 4- position of the benzoyl ring. Other interesting *in vivo* hydrolysable esters include, for example, R<sup>A</sup>C(O)O(C<sub>1-6</sub>)alkyl-CO-, wherein R<sup>A</sup> is for example, benzyloxy-(C<sub>1-4</sub>)alkyl, or phenyl). Suitable substituents on a
- 10 phenyl group in such esters include, for example, 4-(C<sub>1-4</sub>)piperazino-(C<sub>1-4</sub>)alkyl, piperazino-(C<sub>1-4</sub>)alkyl and morpholino-(C<sub>1-4</sub>)alkyl.

- In this specification the generic term "alkyl" includes both straight-chain and branched-chain alkyl groups. However references to individual alkyl groups such as "propyl" are specific for the straight chain version only and references to individual branched-chain
- 15 alkyl groups such as *tert*-butyl are specific for the branched chain version only. For example, "C<sub>1-3</sub>alkyl" includes methyl, ethyl, propyl and isopropyl, examples of "C<sub>1-4</sub>alkyl" include the examples of "C<sub>1-3</sub>alkyl" and butyl and *tert*-butyl and examples of "C<sub>1-6</sub>alkyl" include the examples of "C<sub>1-4</sub>alkyl" and additionally pentyl, 2,3-dimethylpropyl, 3-methylbutyl and hexyl. An analogous convention applies to other generic terms, for example "C<sub>2-4</sub>alkenyl" includes
- 20 vinyl, allyl and 1-propenyl and examples of "C<sub>2-6</sub>alkenyl" include the examples of "C<sub>2-4</sub>alkenyl" and additionally 1-butenyl, 2-butenyl, 3-butenyl, 2-methylbut-2-enyl, 3-methylbut-1-enyl, 1-pentenyl, 3-pentenyl and 4-hexenyl. Examples of "C<sub>2-4</sub>alkynyl" includes ethynyl, 1-propynyl, 2-propynyl and 3-butylnyl and examples of "C<sub>2-6</sub>alkynyl" include the examples of "C<sub>2-4</sub>alkynyl" and additionally 2-pentylnyl, hexynyl and 1-methylpent-2-ynyl.
- 25 Where examples are given of generic terms, these examples are not limiting.

- "Cycloalkyl" is a monocyclic, saturated alkyl ring. The term "C<sub>3-4</sub>cycloalkyl" includes cyclopropyl and cyclobutyl. The term "C<sub>3-5</sub>cycloalkyl" includes "C<sub>3-4</sub>cycloalkyl" and cyclopentyl. The term "C<sub>3-6</sub>cycloalkyl" includes "C<sub>3-5</sub>cycloalkyl" and cyclohexyl. The term "C<sub>3-7</sub>cycloalkyl" includes "C<sub>3-6</sub>cycloalkyl" and additionally cycloheptyl. The term "C<sub>3-</sub>
- 30 10cycloalkyl" includes "C<sub>3-7</sub>cycloalkyl" and additionally cyclooctyl, cyclononyl and cyclodecyl.

"Cycloalkenyl" is a monocyclic ring containing 1, 2, 3 or 4 double bonds. Examples of "C<sub>3-7</sub>cycloalkenyl", "C<sub>5-7</sub>cycloalkenyl" and "C<sub>5-6</sub>cycloalkenyl" are cyclopentenyl, cyclohexenyl and cyclohexadiene and examples of "C<sub>5-10</sub>cycloalkenyl" include these examples and cyclooctatriene.

- 5 Unless otherwise specified "aryl" is monocyclic or bicyclic. Examples of "aryl" therefore include phenyl (an example of monocyclic aryl) and naphthyl (an example of bicyclic aryl).

Examples of "arylC<sub>1-4</sub>alkyl" are benzyl, phenethyl, naphthylmethyl and naphthylethyl.

- Unless otherwise specified "heteroaryl" is a monocyclic or bicyclic aryl ring  
10 containing 5 to 10 ring atoms of which 1, 2, 3 or 4 ring atoms are chosen from nitrogen, sulphur or oxygen where a ring nitrogen or sulphur may be oxidised. Examples of heteroaryl are pyridyl, imidazolyl, quinolinyl, cinnolyl, pyrimidinyl, thienyl, pyrrolyl, pyrazolyl, thiazolyl, oxazolyl, isoxazolyl, pyrazinyl, pyridoimidazolyl, benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl,  
15 indazolyl, indoliziny, isobenzofuranyl, quinazolinyl, imidazopyridinyl and pyrazolopyridinyl. Preferably heteroaryl is pyridyl, imidazolyl, quinolinyl, pyrimidinyl, thienyl, pyrazolyl, thiazolyl, oxazolyl and isoxazolyl. More preferably heteroaryl is pyridyl, imidazolyl and pyrimidinyl. Examples of "monocyclic heteroaryl" are pyridyl, imidazolyl, pyrimidinyl, thienyl, pyrrolyl, pyrazolyl, thiazolyl, oxazolyl, isoxazolyl and pyrazinyl. Examples of  
20 "bicyclic heteroaryl" are quinolinyl, quinazolinyl, cinnolyl, pyridoimidazolyl, benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazolinyl, imidazopyridinyl and pyrazolopyridinyl. Preferred examples B when B is heteroaryl are those examples of bicyclic heteroaryl.

- 25 Examples of "heteroarylC<sub>1-4</sub>alkyl" are pyridylmethyl, pyridylethyl, pyrimidinylethyl, pyrimidinylpropyl, pyrimidinylbutyl, imidazolylpropyl, imidazolylbutyl, quinolinylpropyl, 1,3,4-triazolylpropyl and oxazolylmethyl.

- "Heterocyclyl" is a saturated, partially saturated or unsaturated, monocyclic or bicyclic ring (unless otherwise stated) containing 4 to 12 atoms of which 1, 2, 3 or 4 ring atoms are  
30 chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH<sub>2</sub>- group can optionally be replaced by a -C(O)-; and where unless stated to the contrary a ring nitrogen or sulphur atom is optionally oxidised to form the

N-oxide or S-oxide(s); a ring -NH is optionally substituted by acetyl, formyl, methyl or mesyl; and a ring is optionally substituted by one or more halo. Examples and suitable values of the term "heterocyclyl" are piperidinyl, *N*-acetylpiperidinyl, *N*-methylpiperidinyl, *N*-formylpiperazinyl, *N*-mesylpiperazinyl, homopiperazinyl, piperazinyl, azetidiny, oxetanyl, 5 morpholinyl, tetrahydroisoquinoliny, tetrahydroquinoliny, indoliny, pyranyl, dihydro-2*H*-pyranyl, tetrahydrofuranyl, 2,5-dioximidazolidiny, 2,2-dimethyl-1,3-dioxolanyl and 3,4-methylenedioxyphenyl. Preferred values are 3,4-dihydro-2*H*-pyran-5-yl, tetrahydrofuran-2-yl, 2,5-dioximidazolidiny, 2,2-dimethyl-1,3-dioxolan-2-yl, 2,3-methylenedioxyphenyl and 3,4-methylenedioxyphenyl. Other values are pyridoimidazolyl, benzimidazolyl, benzofuranyl, 10 benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazoliny, imidazopyridiny, pyrazolopyridiny, indoliny, tetrahydroquinoline, tetrahydroisoquinoline and isoindoliny. Examples of monocyclic heterocyclyl are piperidinyl, *N*-acetylpiperidinyl, *N*-methylpiperidinyl, *N*-formylpiperazinyl, *N*-mesylpiperazinyl, homopiperazinyl, piperazinyl, azetidiny, oxetanyl, 15 morpholinyl, pyranyl, tetrahydrofuranyl, 2,5-dioximidazolidiny and 2,2-dimethyl-1,3-dioxolanyl. Examples of bicyclic heterocyclyl are pyridoimidazolyl, benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazoliny, imidazopyridiny, pyrazolopyridiny, indoliny, tetrahydroquinoliny, tetrahydroisoquinoliny, isoindoliny, 2,3-20 methylenedioxyphenyl, and 3,4-dimethylenedioxyphenyl. Examples of saturated heterocyclyl are piperidinyl, pyrrolidinyl and morpholinyl.

The term "halo" refers to fluoro, chloro, bromo and iodo.

Examples of "C<sub>1-3</sub>alkoxy" and "C<sub>1-4</sub>alkoxy" include methoxy, ethoxy, propoxy and isopropoxy. Examples of "C<sub>1-6</sub>alkoxy" include the examples of "C<sub>1-4</sub>alkoxy" and additionally 25 pentyloxy, 1-ethylpropoxy and hexyloxy.

"Heteroalkyl" is alkyl containing at least one carbon atom and having at least one carbon atom replaced by a hetero group independently selected from N, O, S, SO, SO<sub>2</sub>, (a hetero group being a hetero atom or group of atoms). Examples include -CH<sub>2</sub>O-, OCH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>O-, -CH<sub>2</sub>SCH<sub>2</sub>CH<sub>2</sub> and -OCH(CH<sub>3</sub>)<sub>2</sub>-.

30 "HaloC<sub>1-4</sub>alkyl" is a C<sub>1-4</sub>alkyl group substituted by one or more halo. Examples of "haloC<sub>1-4</sub>alkyl" include fluoromethyl, trifluoromethyl, 1-chloroethyl, 2-chloroethyl, 2-

bromopropyl, 1-fluoroisopropyl and 4-chlorobutyl. Examples of "haloC<sub>1-6</sub>alkyl" include the examples of "haloC<sub>1-4</sub>alkyl" and 1-chloropentyl, 3-chloropentyl and 2-fluorohexyl.

Examples of "hydroxyC<sub>1-4</sub>alkyl" include hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxypropyl, 1-hydroxyisopropyl and 4-hydroxybutyl.

5 Example of "C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl" include methoxymethyl, ethoxymethyl, methoxyethyl, methoxypropyl and propoxybutyl.

"HaloC<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl" is a C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl group substituted by one or more halo. Examples of "haloC<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl" include trifluoromethoxymethyl, 1-(chloromethoxy)ethyl, 2-fluoroethoxymethyl, 2-(4-bromobutoxy)ethyl and 2-(2-

10 iodoethoxy)ethyl.

Examples of "carboxyC<sub>1-4</sub>alkyl" include carboxymethyl, 2-carboxyethyl and 2-carboxypropyl.

A "carbocyclic 5 to 6-membered" ring is (unless specifically stated) a saturated, partially saturated or unsaturated ring containing 5 to 6 ring carbon atoms. Examples include 15 cyclopentyl, cyclopent-3-enyl, cyclohexyl and cyclopent-2-enyl. An analogous convention applies for a "carbocyclic 3 to 7-membered" ring which includes the examples a "carbocyclic 5 to 6-membered" ring and additionally cyclopropyl and cyclobutyl.

Heterocyclic rings are rings containing 1, 2 or 3 ring atoms selected from nitrogen, oxygen and sulphur. "Heterocyclic 4 to 6-membered", "heterocyclic 5 to 6-membered" and 20 "heterocyclic 5 to 7-membered" rings are pyrrolidinyl, piperidinyl, piperazinyl, homopiperidinyl, homopiperazinyl, thiomorpholinyl, thiopyranyl and morpholinyl. "Heterocyclic 4 to 7-membered" rings include the examples of "heterocyclic 5 to 7-membered" and additionally azetidyl. Saturated heterocyclic 3- to 7-membered, 4- to 7-membered and 5- to 6-membered rings include piperidinyl, pyrrolidinyl and morpholinyl.

25 Where optional substituents are chosen from "one or more" groups or substituents it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups. Preferably "one or more" means "1, 2 or 3" and this is particularly the case when the group or substituent is halo. "One or more" may also means "1 or 2".

30 Compounds of the present invention have been named with the aid of computer software (ACD/Name version 5.09).

Preferred values of Y<sup>1</sup>, Y<sup>2</sup>, z, n, W, m, D, X, B, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are as follows.

Such values may be used where appropriate with any of the definitions, claims or embodiments defined herein.

In one aspect of the invention  $Y^1$  and  $Y^2$  are both O.

In one aspect of the invention  $z$  is  $NR^8$ .

5 In one aspect of the invention  $n$  is 1. In another aspect  $n$  is 0.

In one aspect of the invention  $W$  is  $NR^1$ . In another aspect  $W$  is  $CR^1R^2$ . In a further aspect  $W$  is a bond.

In one aspect of the invention  $m$  is 0. In another aspect  $m$  is 1.

In one aspect of the invention  $D$  is hydrogen, methyl or fluoro. In another aspect  $D$  is  
10 hydrogen.

In one aspect of the invention  $X$  is  $-CR^{12}R^{13}-Q-$  or  $-CR^{12}R^{13}-Q-CR^{14}R^{15}-$ . In another aspect of the invention  $X$  is  $-CR^{12}R^{13}-Q-$ ,  $-Q-CR^{14}R^{15}-$  or  $-CR^{12}R^{13}-Q-CR^{14}R^{15}-$ . In another aspect  $X$  is  $Q$ . In a further aspect  $X$  is  $-(CH_2)-O-$ ,  $-O-(CH_2)-$ ,  $-(CH_2)-O-(CH_2)-$  or  $-(CHMe)-O-$  or  $O$ . In yet another aspect  $X$  is  $-(CH_2)-O-$  or  $-O-(CH_2)-$

15 In one aspect of the invention  $Q$  is O.

In one aspect, when  $n$  is 1 and  $W$  is  $NR^1$ ,  $CR^1R^2$  or a bond; or when  $n$  is 0 and  $W$  is  $CR^1R^2$ ;  $B$  is a group selected from aryl, heteroaryl and heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo,  $C_{1-4}$ alkyl (optionally substituted by one or more halo),  
20  $C_{2-4}$ alkynyl, heteroaryl,  $-OR^9$ , cyano,  $-NR^9R^{10}$ ,  $-CONR^9R^{10}$  and  $-NR^9COR^{10}$ ; or  $B$  is  $C_{2-4}$ alkenyl or  $C_{2-4}$ alkynyl optionally substituted by  $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl or heterocyclyl. In another aspect, when  $n$  is 1 and  $W$  is  $NR^1$ ,  $CR^1R^2$  or a bond; or when  $n$  is 0 and  $W$  is  $CR^1R^2$ ;  $B$  is phenyl, naphthyl, pyridyl, quinoliny, isoquinoliny, thienopyridyl, naphthyridinyl, 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, thienopyrimidinyl, pyridoimidazolyl,  
25 benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazoliny, imidazopyridinyl, pyrazolopyridinyl, indoliny, tetrahydroquinoliny, tetrahydroisoquinoliny or isoindoliny, where each is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo,  $C_{1-4}$ alkyl (optionally substituted  
30 by one or more halo),  $C_{2-4}$ alkynyl, heteroaryl,  $-OR^9$ , cyano,  $-NR^9R^{10}$ ,  $-CONR^9R^{10}$  and  $-NR^9COR^{10}$ ; or  $B$  is vinyl or ethynyl optionally substituted by  $C_{1-4}$ alkyl. In another aspect when  $n$  is 1 and  $W$  is  $NR^1$ ,  $CR^1R^2$  or a bond; or when  $n$  is 0 and  $W$  is  $CR^1R^2$ ;  $B$  is phenyl,

- naphthyl, pyridyl, quinolinyl, isoquinolinyl, thienopyridyl, naphthyridinyl, 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, thienopyrimidinyl, pyridoimidazolyl, benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazolinyl,
- 5 imidazopyridinyl, pyrazolopyridinyl, indolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl or isoindolinyl, where each is optionally substituted by one or more groups independently selected from trifluoromethyl, trifluoromethoxy, fluoro, chloro, bromo, methyl, isopropyl, ethynyl, cyano, acetamido, propyloxy, isopropylloxymethoxy, nitro, pyrrolidinylcarbonyl, *N*-propylcarbamoyl, pyrrolidinyl, piperidinyl, isoxazolyl, pyrazolyl, imidazolyl, oxazolyl,
- 10 thiazolyl, pyrimidinyl and pyridyl; or B is vinyl or ethynyl optionally substituted by methyl or ethyl. In a further aspect when *n* is 1 and W is  $\text{NR}^1$ ,  $\text{CR}^1\text{R}^2$  or a bond; or when *n* is 0 and W is  $\text{CR}^1\text{R}^2$ ; B is quinolin-4-yl, naphthyl, 2-methylquinolin-4-yl, 3-methylnaphthyl, 7-methylquinolin-5-yl, 6-methylquinolin-8-yl, 7-methylisoquinolin-5-yl, 6-methylthieno[2,3-*b*]pyridyl, 5-methylthieno[3,2-*b*]pyridyl, 2-methyl-1,8-naphthyridinyl, 2-
- 15 trifluoromethylquinolin-4-yl, 2-ethynylquinolin-4-yl, 7-chloroquinolin-5-yl, 7-fluoro-2-methylquinolin-4-yl, 2-methyl-*N*-oxoquinolin-4-yl, 3-methylisoquinolin-1-yl, 5-fluoro-2-methylquinolin-4-yl, 2,6-dimethylpyrid-4-yl, 2,5-dimethylpyridin-4-yl, 2,5-dimethylphenyl, 2,5-difluorophenyl, 3,5-difluorophenyl, 2,6-difluoro-3-methylphenyl, 2-chloro-6-fluorophenyl, 3-fluoro-6-methylphenyl, 2,6-difluorophenyl, 3,4-dichlorophenyl, 2-fluoro-3-methylphenyl,
- 20 2,4-dichlorophenyl, 2,6-dichlorophenyl, 2,4,6-trimethylphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 3-chloro-4-methylphenyl, 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, 5-fluoro-2-methylpyridinyl, 2,4-dimethylphenyl, 1-methylquinolinyl, 2-chloro-4-fluorophenyl, 2-chloro-4-trifluoromethylphenyl, 2-bromo-4,6-difluorophenyl, 2-bromo-4-fluorophenyl, 2,4-dichlorophenyl, 2-bromo-4-chlorophenyl, 2-methoxy-4-
- 25 methylphenyl, 4-chloro-2-nitrophenyl, 4-methyl-2-nitrophenyl, 2,4-difluorophenyl, 4-bromo-2-fluorophenyl, 2-methoxy-4-nitrophenyl, 2-chloro-4-nitrophenyl, 4-bromo-2-methoxyphenyl, 2-fluoro-4-nitrophenyl, 2-chloro-4-bromophenyl, 2-chloro-4-methylphenyl, 2-chloro-4-methoxyphenyl, 4-fluoro-2-methoxyphenyl, 2-fluoro-4-chlorophenyl, 4-fluoro-2-methylphenyl, 7-chloroquinolin-4-yl, 8-chloroquinolin-4-yl, 3-chloro-5-trifluoromethylpyrid-
- 30 2-yl, 3,5-dichloropyrid-2-yl, 6-chloroquinolin-4-yl, 5-methylthieno[2,3-*d*]pyrimidin-4-yl, 7-methylthieno[3,2-*d*]pyrimidin-4-yl, 8-fluoroquinolin-4-yl, 4-chloro-2-(isoxazol-5-yl)phenyl, 2-(isoxazol-5-yl)-4-trifluoromethylphenyl, 6-fluoroquinolin-4-yl, 2-methylquinolin-4-yl, 6-



- chloro-2-methylquinolin-4-yl, 1,6-naphthyridin-4-yl, thieno[3,2-*b*]pyrid-7-yl, 5-fluoro-2-(isoxazol-5-yl)phenyl, 4-fluoro-2-(isoxazol-5-yl)phenyl, 4-chloro-2-trifluoromethylphenyl, 2-chloro-5-fluorophenyl, vinyl, ethynyl, prop-1-enyl, prop-1-ynyl or but-1-ynyl. In one aspect, B is a group selected from bicyclic aryl, bicyclic heteroaryl and bicyclic heterocyclyl, where
- 5 each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo, C<sub>1-4</sub>alkyl (optionally substituted by one or more halo), C<sub>2-4</sub>alkynyl, heteroaryl, -OR<sup>9</sup>, cyano, -NR<sup>9</sup>R<sup>10</sup>, -CONR<sup>9</sup>R<sup>10</sup> and -NR<sup>9</sup>COR<sup>10</sup>; or B is C<sub>2-4</sub>alkenyl or C<sub>2-4</sub>alkynyl optionally substituted by C<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl or heterocyclyl. In another aspect B is naphthyl, quinolinyl, isoquinolinyl, thienopyridyl, 2,3-
- 10 methylenedioxyphenyl, 3,4-methylenedioxyphenyl, naphthyridinyl, thienopyrimidinyl, pyridoimidazolyl, benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazolinyl, imidazopyridinyl, pyrazolopyridinyl, indoliny, tetrahydroquinolinyl, tetrahydroisoquinolinyl or isoindolinyl, where each is optionally substituted by one or more
- 15 groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo, C<sub>1-4</sub>alkyl (optionally substituted by one or more halo), C<sub>2-4</sub>alkynyl, heteroaryl, -OR<sup>9</sup>, cyano, -NR<sup>9</sup>R<sup>10</sup>, -CONR<sup>9</sup>R<sup>10</sup> and -NR<sup>9</sup>COR<sup>10</sup>; or B is vinyl or ethynyl optionally substituted by C<sub>1-4</sub>alkyl. In another aspect B is naphthyl, quinolinyl, isoquinolinyl, thienopyridyl, 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, naphthyridinyl, thienopyrimidinyl,
- 20 pyridoimidazolyl, benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazolinyl, imidazopyridinyl, pyrazolopyridinyl, indoliny, tetrahydroquinolinyl, tetrahydroisoquinolinyl or isoindolinyl, where each is optionally substituted by one or more groups independently selected from trifluoromethyl, trifluoromethoxy, fluoro, chloro, bromo,
- 25 methyl, isopropyl, ethynyl, cyano, acetamido, propyloxy, isopropyloxy, methoxy, nitro, pyrrolidinylcarbonyl, *N*-propylcarbonyl; or B is vinyl or ethynyl optionally substituted by methyl or ethyl. In another aspect B is quinolin-4-yl, naphthyl, 2-methylquinolin-4-yl, 3-methylnaphthyl, 7-methylquinolin-5-yl, 6-methylquinolin-8-yl, 7-methylisoquinolin-5-yl, 6-methylthieno[2,3-*b*]pyridyl, 5-methylthieno[3,2-*b*]pyridyl, 2-methyl-1,8-naphthyridinyl, 2-
- 30 trifluoromethylquinolin-4-yl, 2-ethynylquinolin-4-yl, 7-chloroquinolin-5-yl, 7-fluoro-2-methylquinolin-4-yl, 2-methyl-*N*-oxoquinolin-4-yl, 3-methylisoquinolin-1-yl, 5-fluoro-2-methylquinolin-4-yl, 3,4-methylenedioxyphenyl, 1-methylquinolinyl, 7-chloroquinolin-4-yl, 8-



chloroquinolin-4-yl, 6-chloroquinolin-4-yl, 5-methylthieno[2,3-*d*]pyrimidin-4-yl, 7-methylthieno[3,2-*d*]pyrimidin-4-yl, 8-fluoroquinolin-4-yl, 6-fluoroquinolin-4-yl, 2-methylquinolin-4-yl, 6-chloro-2-methylquinolin-4-yl, 1,6-naphthyridin-4-yl, thieno[3,2-*b*]pyrid-7-yl, vinyl, ethynyl, prop-1-enyl, prop-1-ynyl or but-1-ynyl. In another aspect B is a group selected from aryl and heteroaryl where each group is optionally substituted by one or more groups independently selected from halo, C<sub>1-4</sub>alkyl (optionally substituted by one or more halo), heteroaryl and C<sub>2-4</sub>alkynyl. In another aspect B is a group selected from quinolinyl, pyridyl and phenyl where each group is optionally substituted by one or more methyl, trifluoromethyl, trifluoromethoxy, halo or isoxazolyl. In a further aspect B is 2-methylquinolin-4-yl, 2,5-dimethylphenyl or 2,5-dimethylpyrid-4-yl. In yet another aspect B is 2-methylquinolin-4-yl.

In one aspect of the invention R<sup>1</sup> is hydrogen or methyl. In another aspect R<sup>1</sup> is hydrogen.

In one aspect of the invention R<sup>2</sup> is hydrogen or methyl. In another aspect R<sup>2</sup> is hydrogen.

In one aspect of the invention R<sup>3</sup> is hydrogen, methyl, ethyl, propyl or phenyl. In another aspect R<sup>3</sup> is hydrogen.

In one aspect of the invention R<sup>4</sup> is hydrogen or methyl. In another aspect R<sup>4</sup> is hydrogen.

In one aspect of the invention R<sup>5</sup> is hydrogen or methyl. In another aspect R<sup>5</sup> is hydrogen.

In one aspect of the invention R<sup>6</sup> is hydrogen or methyl. In another aspect R<sup>6</sup> is hydrogen.

In one aspect of the invention R<sup>1</sup> and R<sup>3</sup> together with the nitrogen or carbon and carbon to which they are respectively attached form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and SO<sub>2</sub> where the ring is optionally substituted by one or more C<sub>1-4</sub>alkyl. In another aspect R<sup>1</sup> and R<sup>3</sup> together with the nitrogen or carbon and carbon to which they are respectively attached form a piperidine, pyrrolidine, piperazine, morpholine, cyclohexane or cyclopentane ring.

In one aspect of the invention R<sup>3</sup> and R<sup>4</sup> together form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and SO<sub>2</sub> where the ring is optionally substituted by one or more C<sub>1-4</sub>alkyl.

In one aspect of the invention  $R^5$  and  $R^6$  together form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted by one or more  $C_{1-4}$ alkyl.

In one aspect of the invention  $R^7$  is hydrogen or a group selected from  $C_{1-4}$ alkyl,  $C_{3-5}$ cycloalkyl, aryl, heteroaryl or heterocyclyl where the group is optionally substituted by heterocyclyl, aryl and heteroaryl; and wherein the group from which  $R^7$  may be selected is optionally substituted on the group and/or on its optional substituent by one or more substituents independently selected from halo, cyano,  $C_{1-4}$ alkyl,  $-OR^{21}$ ,  $-CO_2R^{21}$ ,  $-NR^{21}COR^{22}$ ,  $-NR^{21}CO_2R^{22}$  and  $-CONR^{21}R^{22}$ . In one aspect  $R^7$  is hydrogen or a group selected from  $C_{1-4}$ alkyl,  $C_{3-5}$ cycloalkyl, aryl, heteroaryl or heterocyclyl where the group is optionally substituted by heterocyclyl, aryl and heteroaryl; and wherein the group from which  $R^7$  may be selected is optionally substituted on the group and/or on its optional substituent by one or more substituents independently selected from halo, cyano,  $C_{1-4}$ alkyl,  $-OR^{21}$ ,  $-CO_2R^{21}$ , and  $-NR^{21}CO_2R^{22}$ . In another aspect  $R^7$  is hydrogen or a group selected from  $C_{1-4}$ alkyl, aryl,  $C_{1-4}$ alkyl, heteroaryl,  $C_{1-4}$ alkyl, heterocyclyl,  $C_{1-4}$ alkyl, aryl, heteroaryl, heterocyclyl and  $C_{3-5}$ cycloalkyl where the group is optionally substituted by cyano,  $C_{1-4}$ alkyl, halo,  $-OR^{21}$ ,  $-NR^{21}R^{22}$ ,  $-CO_2R^{21}$  and  $-NR^{21}CO_2R^{22}$ . In a further aspect  $R^7$  is selected from hydrogen, methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, *tert*-butyl, isobutyl, 1-hydroxyethyl, 2-hydroxyethyl, 3-hydroxypropyl, methoxymethyl, 2-methoxyethyl, 2-cyanoethyl, 2-aminoethyl, phenyl, pyridyl, benzyl, 3-methylbenzyl, phenylethyl, 4-chlorophenylethyl, 4-fluorophenylethyl, phenylpropyl, 4-chlorophenylpropyl, 4-fluorophenylpropyl, 4-methylpiperazin-1-ylethyl, morpholin-4-ylpropyl, pyrimidin-2-ylethyl, pyrimidin-2-ylpropyl, pyrimidin-2-ylbutyl, 5-fluoropyrimidin-2-ylpropyl, imidazol-1-ylpropyl, imidazol-1-ylbutyl, 1,3,4-triazolylpropyl, piperidiny, tetrahydro-2H-pyranyl, tetrahydro-2H-pyranylmethyl, pyrid-2-ylmethyl, pyrid-4-ylmethyl, pyrid-3-ylmethyl, piperidin-4-ylmethyl, N-(*tert*-butoxycarbonyl)piperidin-4-yl, *tert*-butoxycarbonylaminomethyl, N-(methylcarbonyl)piperidin-4-yl, benzyloxyethyl, N-(*tert*-butoxycarbonyl)piperidin-4-ylmethyl, (3,4,4-trimethyl-2,5-dioximidazolidin-1-yl)methyl and N-benzoyl-N-phenylaminomethyl. In a further aspect  $R^7$  is hydrogen or  $C_{1-4}$ alkyl optionally substituted with halo, hydroxy or  $C_{1-3}$ alkoxy. In yet another aspect  $R^7$  is hydrogen, methyl or ethyl.

In one aspect of the invention  $R^3$  and  $R^7$  together with the carbon atoms to which they are each attached and  $(CR^5R^6)_n$  form a saturated 5- to 7-membered ring optionally containing

a heteroatom group selected from NH, O, S, SO and SO<sub>2</sub> where the ring is optionally substituted on carbon or nitrogen by one or more C<sub>1-4</sub>alkyl. In another aspect R<sup>3</sup> and R<sup>7</sup> together with the carbon atoms to which they are each attached and (CR<sup>5</sup>R<sup>6</sup>)<sub>n</sub> form a piperidinyl, pyrrolidinyl, piperazine, morpholine, cyclohexane or cyclopentane ring.

5 In one aspect of the invention R<sup>8</sup> is hydrogen or methyl. In another aspect R<sup>8</sup> is hydrogen.

In one aspect R<sup>9</sup> is hydrogen or methyl.

In one aspect R<sup>10</sup> is hydrogen or methyl.

In one aspect R<sup>11</sup> is methyl.

10 In one aspect R<sup>12</sup> is hydrogen or methyl.

In one aspect R<sup>13</sup> is hydrogen or methyl.

In one aspect R<sup>14</sup> is hydrogen or methyl.

In one aspect R<sup>15</sup> is hydrogen or methyl.

In one aspect R<sup>16</sup> is hydrogen or methyl.

15 In one aspect R<sup>17</sup> is selected from fluoro, chloro, methyl or methoxy.

In one aspect of the invention R<sup>19</sup> is a group selected from C<sub>1-6</sub>alkyl, aryl and arylC<sub>1-4</sub>alkyl which group is optionally substituted by halo. In another aspect R<sup>19</sup> is a group selected from methyl, phenyl and benzyl which group is optionally substituted by chloro. In one aspect of the invention R<sup>19</sup> is methyl.

20 In one aspect of the invention R<sup>18</sup> is hydrogen or a group selected from C<sub>1-6</sub>alkyl, aryl and arylC<sub>1-4</sub>alkyl which group is optionally substituted by halo. In another aspect R<sup>18</sup> is hydrogen or a group selected from methyl, phenyl and benzyl which group is optionally substituted by chloro.

In one aspect R<sup>20</sup> is hydrogen or methyl.

25 In one aspect R<sup>21</sup> is hydrogen, methyl, ethyl, phenyl and benzyl.

In one aspect R<sup>22</sup> is hydrogen, methyl, ethyl, *tert*-butyl, phenyl and benzyl. In another aspect R<sup>22</sup> is hydrogen or methyl.

In one aspect of the invention R<sup>21</sup> and R<sup>22</sup> are independently hydrogen, C<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkyl, aryl, arylC<sub>1-4</sub>alkyl or benzoyl.

30 In one aspect of the invention R<sup>25</sup> is a group selected from C<sub>1-6</sub>alkyl, aryl and arylC<sub>1-4</sub>alkyl which group is optionally substituted by halo. In another aspect R<sup>25</sup> is a group selected

from methyl, phenyl and benzyl which group is optionally substituted by chloro. In one aspect of the invention  $R^{25}$  is methyl.

A preferred class of compound is of formula (1) wherein:

- 5  $Y^1$  and  $Y^2$  are both O;  
 $z$  is  $NR^8$ ;  
 $n$  is 1 and  $W$  is  $NR^1$ ,  $CR^1R^2$  or a bond; or  $n$  is 0 and  $W$  is  $CR^1R^2$ ;  
 $m$  is 1;  
 $D$  is hydrogen, methyl or fluoro;
- 10  $X$  is  $-CR^{12}R^{13}-Q-$ ,  $-Q-CR^{14}R^{15}-$ ,  $-CR^{12}R^{13}-Q-CR^{14}R^{15}-$  or  $Q$ ;  
 $Q$  is O;  
 $B$  is a group selected from aryl, heteroaryl and heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo, cyano,  $C_{1-4}$ alkyl (optionally substituted one or more halo),  $C_{2-4}$ alkenyl, heteroaryl,  $-OR^9$ ,  $-NR^9R^{10}$ ,  $-CONR^9R^{10}$  and  $-NR^9COR^{10}$ ; or  $B$  is  $C_{2-4}$ alkenyl or  $C_{2-4}$ alkynyl, optionally substituted by  $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl or heterocyclyl;  
 $R^1$  and  $R^2$  are independently hydrogen or methyl;  
 $R^3$  is hydrogen, methyl, ethyl, propyl or phenyl;  
 $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$  and  $R^{15}$  are independently hydrogen or methyl;
- 20  $R^7$  is hydrogen or a group selected from  $C_{1-4}$ alkyl,  $C_{3-5}$ cycloalkyl, aryl, heteroaryl or heterocyclyl where the group is optionally substituted by heterocyclyl, aryl and heteroaryl; and wherein the group from which  $R^7$  may be selected is optionally substituted on the group and/or on its optional substituent by one or more substituents independently selected from halo, cyano,  $C_{1-4}$ alkyl,  $-OR^{21}$ ,  $-CO_2R^{21}$ ,  $-NR^{21}COR^{22}$ ,  $-NR^{21}CO_2R^{22}$  and  $-CONR^{21}R^{22}$ ;
- 25  $R^{21}$  is hydrogen, methyl, ethyl, phenyl or benzyl;  
 $R^{22}$  is hydrogen, methyl, ethyl, *tert*-butyl, phenyl or benzyl.

Another preferred class of compound is of formula (1) wherein:

- $Y^1$  and  $Y^2$  are both O;  
 $z$  is  $NR^8$ ;
- 30  $n$  is 1 and  $W$  is  $NR^1$ ,  $CR^1R^2$  or a bond; or  $n$  is 0 and  $W$  is  $CR^1R^2$ ;  
 $m$  is 1;  
 $D$  is hydrogen, methyl or fluoro;

X is  $-\text{CR}^{12}\text{R}^{13}-\text{Q}-$ ,  $-\text{Q}-\text{CR}^{14}\text{R}^{15}-$ ,  $-\text{CR}^{12}\text{R}^{13}-\text{Q}-\text{CR}^{14}\text{R}^{15}-$  or Q;

Q is O;

B is phenyl, naphthyl, pyridyl, quinoliny, isoquinoliny, thienopyridyl, naphthyridinyl, 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, thienopyrimidinyl, pyridoimidazolyl,

- 5 benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazoliny, imidazopyridinyl, pyrazolopyridinyl, indoliny, tetrahydroquinoliny, tetrahydroisoquinoliny or isoindoliny, where each is optionally substituted by one or more groups independently selected from trifluoromethyl, trifluoromethoxy, fluoro, chloro, bromo, methyl, isopropyl,
- 10 ethynyl, cyano, acetamido, propyloxy, isopropylloxymethoxy, nitro, pyrrolidinylcarbonyl, *N*-propylcarbonyl, pyrrolidinyl, piperidinyl, isoxazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, pyrimidinyl and pyridyl; or B is vinyl or ethynyl optionally substituted by methyl or ethyl.

$\text{R}^1$  and  $\text{R}^2$  are independently hydrogen or methyl;

- 15  $\text{R}^3$  is hydrogen, methyl, ethyl, propyl or phenyl;

$\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^8$ ,  $\text{R}^{12}$ ,  $\text{R}^{13}$ ,  $\text{R}^{14}$  and  $\text{R}^{15}$  are independently hydrogen or methyl;

$\text{R}^7$  is hydrogen or a group selected from  $\text{C}_{1-4}$ alkyl, aryl $\text{C}_{1-4}$ alkyl, heteroaryl $\text{C}_{1-4}$ alkyl, heterocyclyl $\text{C}_{1-4}$ alkyl, aryl, heteroaryl, heterocyclyl and  $\text{C}_{3-5}$ cycloalkyl where the group is optionally substituted by cyano,  $\text{C}_{1-4}$ alkyl, halo,  $-\text{OR}^{21}$ ,  $-\text{NR}^{21}\text{R}^{22}$ ,  $-\text{CO}_2\text{R}^{21}$  and -

- 20  $\text{NR}^{21}\text{CO}_2\text{R}^{22}$ ;

$\text{R}^{21}$  is hydrogen, methyl, ethyl, phenyl and benzyl;

$\text{R}^{22}$  is hydrogen, methyl, ethyl, phenyl, *tert*-butyl and benzyl.

Another preferred class of compound is of formula (1) wherein:

$\text{Y}^1$  and  $\text{Y}^2$  are both O;

- 25 z is  $\text{NR}^8$ ;

n is 0;

W is  $\text{NR}^1$ ;

m is 1;

D is hydrogen, methyl or fluoro;

- 30 X is  $-\text{CR}^{12}\text{R}^{13}-\text{Q}-$ ,  $-\text{Q}-\text{CR}^{14}\text{R}^{15}-$ ,  $-\text{CR}^{12}\text{R}^{13}-\text{Q}-\text{CR}^{14}\text{R}^{15}-$  or Q;

Q is O;

B is a group selected from bicyclic aryl, bicyclic heteroaryl and bicyclic heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo, cyano, C<sub>1-4</sub>alkyl (optionally substituted by one or more halo), C<sub>2-4</sub>alkenyl, heteroaryl, -OR<sup>9</sup>, -NR<sup>9</sup>R<sup>10</sup>, -CONR<sup>9</sup>R<sup>10</sup> and -NR<sup>9</sup>COR<sup>10</sup>; or B is C<sub>2-</sub>

5 <sub>4</sub>alkenyl or C<sub>2-4</sub>alkynyl, optionally substituted by C<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl or heterocyclyl;

R<sup>1</sup> is hydrogen;

R<sup>3</sup> is hydrogen, methyl, ethyl, propyl or phenyl;

R<sup>4</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> are independently hydrogen or methyl;

R<sup>7</sup> is hydrogen or a group selected from C<sub>1-4</sub>alkyl, arylC<sub>1-4</sub>alkyl, heteroarylC<sub>1-4</sub>alkyl,

10 heterocyclylC<sub>1-4</sub>alkyl, aryl, heteroaryl, heterocyclyl and C<sub>3-5</sub>cycloalkyl where the group is optionally substituted by cyano, C<sub>1-4</sub>alkyl, halo, -OR<sup>21</sup>, -NR<sup>21</sup>R<sup>22</sup>, -CO<sub>2</sub>R<sup>21</sup> and -NR<sup>21</sup>CO<sub>2</sub>R<sup>22</sup>;

R<sup>21</sup> is hydrogen, methyl, ethyl, phenyl and benzyl;

R<sup>22</sup> is hydrogen, methyl, ethyl, phenyl, *tert*-butyl and benzyl.

15 Another preferred class of compound is of formula (1) wherein:

Y<sup>1</sup> and Y<sup>2</sup> are both O;

z is NR<sup>8</sup>;

n is 0 or 1;

W is NR<sup>1</sup>, CR<sup>1</sup>R<sup>2</sup> or a bond;

20 m is 1;

D is hydrogen, methyl or fluoro;

X is -CR<sup>12</sup>R<sup>13</sup>-Q-, -Q-CR<sup>14</sup>R<sup>15</sup>- or -CR<sup>12</sup>R<sup>13</sup>-Q-CR<sup>14</sup>R<sup>15</sup>-;

Q is O;

B is 2-methylquinolin-4-yl;

25 R<sup>1</sup> and R<sup>2</sup> are independently hydrogen or methyl;

R<sup>3</sup> is hydrogen, methyl, ethyl or phenyl;

R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>8</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> are independently hydrogen or methyl;

R<sup>7</sup> is hydrogen or C<sub>1-4</sub>alkyl optionally substituted with halo, hydroxy or C<sub>1-3</sub>alkoxy.

In another aspect of the invention, preferred compounds of the invention are any one  
30 of:

5-[(4-[(2,5-dimethylbenzyl)oxy]piperidin-1-yl)sulphonyl)methyl]-5-methylimidazolidine-2,4-dione; and


5-[(4-(2-methylquinolin-4-ylmethoxy)piperidin-1-yl)sulphonylmethyl]-5-methylimidazolidine-2,4-dione.

In another aspect of the invention, preferred compounds are any one of:

- 5 R/S-5-[(4-(2-methylquinolin-4-ylmethoxy)piperidin-1-yl)sulphonylmethyl]-5-methylimidazolidine-2,4-dione;
- 5-[2-((4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)ethyl]imidazolidine-2,4-dione;
- 5-{2-[(4-[(2-methylquinolin-4-yl)oxy]methyl)piperidin-1-yl)sulphonyl]ethyl}imidazolidine-
- 10 2,4-dione;
- 5-methyl-5-[(4-[(2-methylquinolin-4-yl)oxy]methyl)piperidin-1-yl)sulphonylmethyl}imidazolidine-2,4-dione;
- 5-ethyl-5-[(4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonylmethyl}imidazolidine-2,4-dione;
- 15 5-methyl-5-[2-((4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)ethyl]imidazolidine-2,4-dione;
- 5-ethyl-5-[2-((4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)ethyl]imidazolidine-2,4-dione;
- (5S)-5-methyl-5-{4-[(2-methylquinolin-4-yl)methoxymethyl]piperidylsulphonylmethyl}-2,4-
- 20 dioxoimidazolidine; and
- (5S)-5-ethyl-5-{4-[(2-methylquinolin-4-yl)methoxymethyl]piperidylsulphonylmethyl}-2,4-dioxoimidazolidine.

In another aspect the present invention provides a process for the preparation of a compound of formula (1) or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof which comprises:

- a) converting a ketone or aldehyde of formula (2) into a compound of formula (1);



and thereafter if necessary:




- i) converting a compound of formula (1) into another compound of formula (1);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

The hydantoin can be prepared by a number of methods for example;

- 5 a) The aldehyde or ketone may be reacted with ammonium carbonate and potassium cyanide in aqueous alcohols using the method of Bucherer and Berge (*Adv. Het. Chem.*, 1985, 38, 177).
- b) The aldehyde or ketone can be first converted to the cyanohydrin and then further reacted with ammonium carbonate (*Chem. Rev.*, 1950, 56, 403).
- 10 c) The aldehyde or ketone can be converted to the alpha-amino nitrile and then either reacted with ammonium carbonate or aqueous carbon dioxide or potassium cyanate followed by mineral acid (*Chem. Rev.*, 1950, 56, 403).


The process may further comprise a process for the preparation of a ketone or aldehyde of formula (2) where W is a bond and n is 0 (indicated as a compound of formula (2')) which  
 15 process comprises reacting a sulphonamide of formula (3) with a compound of formula (4) where LG represents a leaving group such as halo, alkoxy or aryloxy.



Scheme 2

This process comprises the reaction of the sulphonamide of formula (3) with a base such as  
 20 lithium bis(trimethylsilyl)amide or lithium diisopropylamide in an inert solvent such as tetrahydrofuran at temperatures from -78°C to 0°C for 1 to 2 hours followed by addition of a compound of formula (4) at a temperature of -78°C to room temperature for 1 to 24 hours. A compound of formula (4) is commercially available or can be easily prepared by the skilled man.

25 A ketone of formula (2') may additionally be prepared by the process illustrated in Scheme 3:




Scheme 3

The silyl group present in the compound of formula (30) can be removed by tetrabutylammonium fluoride. Suitable leaving groups (L) are halo, mesyl and tosyl. A

- 5 suitable chlorinating agent is POCl<sub>3</sub>. A compound of formula (2') is prepared in the last stage by reacting the compound of formula (33) with the appropriate piperidine reagent. A compound of formula (28) is commercially available or can be easily prepared by the skilled person.

- Alternatively a process for the preparation of a ketone or aldehyde of formula (2) where W is a bond and n is 1 (indicated as a compound of formula (2'')) comprises reacting a sulphonamide of formula (3) with a compound of formula (5) (an epoxide or equivalent) to give an alcohol of formula (6) and oxidising the alcohol to give a ketone or aldehyde of formula (2''):




## Scheme 4

More specifically the process of Scheme 4 comprises the steps of:

- a) reacting the sulphonamide of formula (3) with a base such as lithium diisopropylamide or lithium bis(trimethylsilyl)amide in tetrahydrofuran at a temperature of  $-78^{\circ}\text{C}$  to  $0^{\circ}\text{C}$  for 1 to 2 hours followed by addition of an epoxide or equivalent of formula (5) and reaction for 1 to 24 hours at a temperature of  $-78^{\circ}\text{C}$  to room temperature to give an alcohol of formula (6); and
  - b) oxidation of an alcohol of formula (6) to a ketone or aldehyde of formula (2''), suitable reagents are manganese dioxide, pyridinium chlorochromate, pyridinium dichromate or dimethyl sulphoxide/oxalyl chloride/triethylamine.
- 10 The epoxide or equivalent of formula (5) is commercially available or can be easily prepared by the skilled person.

In another aspect of the invention there is provided a process for the preparation of a compound of formula (1) where W is  $\text{NR}^1$ ,  $\text{R}^1$  is hydrogen and n is 0 (indicated as a compound of formula (1')) which process comprises reaction of a sulphamoyl chloride

- 15 derivative of formula (7) with an amino-hydantoin derivative of formula (8).




## Scheme 5

- Suitable reaction conditions for such a transformation involve the addition of the sulphamoyl chloride to the amino-hydantoin in an inert solvent such as dichloromethane in the presence of
- 20 a base such as triethylamine, pyridine or N,N-diisopropylethylamine at temperature of  $0^{\circ}\text{C}$  to  $50^{\circ}\text{C}$ .

Also provided is a process for the preparation of a hydantoin of formula (8) as shown in Scheme 6:

-28-




Scheme 6

The process of Scheme 6 comprises the steps of:

- reacting dibenzylamine with a halo ketone or aldehyde (where X is halo) of formula (9) in an inert solvent such as tetrahydrofuran or dichloromethane in the presence of a base e.g. triethylamine at room temperature for 24 hours to give a protected amino ketone or aldehyde of formula (10);
- reacting the ketone or aldehyde under hydantoin formation conditions to give a hydantoin of formula (11); and
- removing the benzyl protecting groups by reaction with palladium/hydrogen to yield a hydantoin of formula (8).

A halo ketone or aldehyde of formula (9) is commercially available or can be prepared easily by the skilled person.

Also provided is a process for the preparation of a sulphamoyl chloride of formula (7) as shown in Scheme 7.




Scheme 7

This reaction involves the treatment of a piperidine of formula (12) with sulphonyl chloride in an inert solvent in the presence of a base such as triethylamine or N,N-diisopropylethylamine.

- A piperidine of formula (12) is commercially available or can be easily prepared by the skilled person.


Also provided is a process for the preparation of a compound of formula (1) where W is  $\text{NR}^1$ ,  $\text{R}^1$  is hydrogen and n is 1 (indicated as a compound of formula (1')) which process comprises reacting a sulphonyl chloride derivative of formula (7) with an amino-hydantoin derivative of formula (13).



Scheme 8

Suitable reaction conditions for such a transformation involve the addition of the sulphonyl chloride to the amino-hydantoin in an inert solvent such as dichloromethane in the presence of a base such as triethylamine, pyridine or N,N-diisopropylethylamine at temperature of 0°C to 10 50°C.

Also provided is a process for the preparation of a hydantoin of formula (13), where  $\text{R}^6$  is hydrogen as shown in Scheme 9:



Scheme 9


15 The process of Scheme 9 comprises the steps of:

- reacting an enone of formula (14) with phthalimide in the presence of sodium methoxide in a polar solvent such as dimethyl sulphoxide to give an N-substituted phthalimide of formula (15);
- forming of the hydantoin of formula (16) using e.g. ammonium carbonate and 20 potassium cyanide in aqueous alcohols; and
- removing the phthalimide residue e.g. by reacting with HCl in acetic acid to yield a hydantoin of formula (13).

An enone of formula (14) is commercially available or can be easily prepared by the skilled person.

In another aspect of the invention, there is provided a process for the preparation of compounds of formula (3) (see Scheme 2 and 4) which process is outlined in Scheme 10 and comprises;

- a) reacting a compound of formula (16) with a compound of formula (17) in the presence of a base to deprotonate the compound of formula (17), to yield a compound of formula (18);
- b) removing the protecting group (PG) from the compound of formula (18) to yield a compound of formula (19); wherein X is  $-(CR^9R^{10})_r-Q-(CR^{11}R^{12})_u-$ ;
- c) reacting the compound of formula (19) with a suitable reagent to yield a compound of formula (3);




Scheme 10

In Scheme 10: L is a suitable leaving group such as halo (chloro, bromo, iodo), mesyl, tosyl; a compound of formula (17) can be deprotonated with a base such as sodium hydride, lithium diisopropylamide, butyllithium, lithium bis(trimethylsilyl)amide and reacted with a compound of formula (16) at temperatures ranging from  $-78^{\circ}\text{C}$  to  $70^{\circ}\text{C}$  in an aprotic solvent, e.g. tetrahydrofuran under argon; suitable protecting groups (PG) include Boc (*tert*-butoxycarbonyl), CBz (carboxybenzyl) groups and mesyl or another alkylsulphonyl; in the case where PG is alkylsulphonyl, reaction of formula (16) and formula (17) directly produce a compound of formula (3); a compound of formula (18) can be converted to a compound of formula (19) by treatment with acid (Boc) or hydrogen/ palladium (CBz); a compound of formula (19) can be converted to a compound of formula (3) by treatment with an alkylsulphonyl chloride in the presence of a base such as pyridine in a solvent such as dichloromethane.

A compound of formula (3) can also be prepared by a process as outlined in Scheme 11, which comprises;

- a) reacting a compound of formula (20) with a compound of formula (21), in the presence of a base to yield a compound of formula (18);
- b) removing the protecting group (PG) from the compound of formula (18) to yield a compound of formula (19);

- c) reacting the compound of formula (19) with a suitable reagent to yield a compound of formula (3); and
- d) oxidising Q as required.



5

Scheme 11

- In both schemes 10 and 11: L is a suitable leaving group such as halo (chloro, bromo, iodo), hydroxy, mesyl, nosyl and tosyl; suitable bases to deprotonate compounds of formula (17) and formula (20) include bases such as caesium fluoride, sodium hydride, lithium diisopropylamide, butyllithium and lithium bis(trimethylsilyl)amide; suitable reaction conditions for step a) are temperatures ranging from  $-78^{\circ}\text{C}$  to  $70^{\circ}\text{C}$  and in aprotic solvent, e.g. tetrahydrofuran under argon; suitable protecting groups (PG) include Boc (*tert*-butoxycarbonyl), CBz (carbonyloxybenzyl) groups and mesyl or another alkylsulphonyl; in the case where PG is alkylsulphonyl, reaction of formula (16) and (17) and of formula (20) and formula (21) directly produces a compound of formula (3); a compound of formula (18) can be converted to a compound formula (19) by treatment with acid (Boc) or hydrogen/palladium (CBz); a compound of formula (19) can be converted to a compound of formula (3) by treatment with an alkylsulphonyl chloride in the presence of a base such as pyridine in a solvent such as dichloromethane; and when B is aromatic, X is O and L is OH, Mitsunobu conditions can be used to form a compound of formula (18), i.e. a compound of formula (16) or formula (20) is reacted with a mixture of diethyl azodicarboxylate or diisopropylazodicarboxylate and triphenylphosphine and formula (17) or formula (21) to give a compound of formula (3).
- Compounds of formula (16), (17), (20) and (21) are commercially available or can be easily prepared by the skilled person.


25

In another aspect of the invention, a compound of formula (1) can be prepared by a process which comprises:

- a) reacting a sulphonyl chloride of formula (22) with a piperidine derivative of formula (19) (see scheme 10 or 11 for its preparation).



-32-




Scheme 12

and thereafter if necessary

- i) converting a compound of formula (1) into another compound of formula (1);
- 5 ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

The sulphonyl chloride of formula (22) may be prepared as shown in scheme 13;



Scheme 13

Compounds of formula (24) are readily available or can be easily made by the skilled person. Details of conditions suitable for hydantoin condition are provided herein (see scheme 1).

- 15 It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a
- 20 substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as
- 25 aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group

using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogen group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *tert*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *tert*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with

a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *tert*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

10 The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

As stated hereinbefore the compounds defined in the present invention possesses metalloproteinases inhibitory activity, and in particular TACE inhibitory activity. This property may be assessed, for example, using the procedure set out below.

#### 15 Isolated Enzyme Assays

##### Matrix Metalloproteinase family including for example MMP13.

Recombinant human proMMP13 may be expressed and purified as described by Knauper *et al.* [V. Knauper *et al.*, (1996) *The Biochemical Journal* **271**:1544-1550 (1996)]. The purified enzyme can be used to monitor inhibitors of activity as follows: purified  
20 proMMP13 is activated using 1mM amino phenyl mercuric acid (APMA) 20 hours at 21°C; the activated MMP13 (11.25ng per assay) is incubated for 4-5 hours at 35°C in assay buffer (0.1M Tris-HCl, pH 7.5 containing 0.1M NaCl, 20mM CaCl<sub>2</sub>, 0.02 mM ZnCl and 0.05% (w/v) Brij 35 using the synthetic substrate 7-methoxycoumarin-4-yl)acetyl.Pro.Leu.Gly.Leu.N-3-(2,4-dinitrophenyl)-L-2,3-diaminopropionyl-Ala.Arg.NH<sub>2</sub> in  
25 the presence or absence of inhibitors. Activity is determined by measuring the fluorescence at  $\lambda_{ex}$  328nm and  $\lambda_{em}$  393nm. Percent inhibition is calculated as follows: % Inhibition is equal to the  $[\text{Fluorescence}_{\text{plus inhibitor}} - \text{Fluorescence}_{\text{background}}]$  divided by the  $[\text{Fluorescence}_{\text{minus inhibitor}} - \text{Fluorescence}_{\text{background}}]$ .

A similar protocol can be used for other expressed and purified pro MMPs using  
30 substrates and buffers conditions optimal for the particular MMP, for instance as described in C. Graham Knight *et al.*, (1992) *FEBS Lett.* **296**(3):263-266.

##### Adamalysin family including for example TNF convertase

The ability of the compounds to inhibit proTNF- $\alpha$  convertase enzyme (TACE) may be assessed using a partially purified, isolated enzyme assay, the enzyme being obtained from the membranes of THP-1 as described by K. M. Mohler *et al.*, (1994) *Nature* 370:218-220. The purified enzyme activity and inhibition thereof is determined by incubating the partially

5 purified enzyme in the presence or absence of test compounds using the substrate 4',5'-Dimethoxy-fluoresceinyl Ser.Pro.Leu.Ala.Gln.Ala.Val.Arg.Ser.Ser.Ser.Arg.Cys(4-(3-succinimid-1-yl)-fluorescein)-NH<sub>2</sub> in assay buffer (50mM Tris HCl, pH 7.4 containing 0.1% (w/v) Triton X-100 and 2mM CaCl<sub>2</sub>), at 26°C for 4 hours. The amount of inhibition is determined as for MMP13 except  $\lambda_{ex}$  485nm and  $\lambda_{em}$  538nm were used. The substrate was

10 synthesised as follows. The peptidic part of the substrate was assembled on Fmoc-NH-Rink-MBHA-polystyrene resin either manually or on an automated peptide synthesiser by standard methods involving the use of Fmoc-amino acids and O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) as coupling agent with at least a 4- or 5-fold excess of Fmoc-amino acid and HBTU. Ser<sup>1</sup> and Pro<sup>2</sup> were double-coupled. The

15 following side chain protection strategy was employed; Ser<sup>1</sup>(But), Gln<sup>5</sup>(Trityl), Arg<sup>8,12</sup>(Pmc or Pbf), Ser<sup>9,10,11</sup>(Trityl), Cys<sup>13</sup>(Trityl). Following assembly, the N-terminal Fmoc-protecting group was removed by treating the Fmoc-peptidyl-resin with in DMF. The amino-peptidyl-resin so obtained was acylated by treatment for 1.5-2hr at 70°C with 1.5-2 equivalents of 4',5'-dimethoxy-fluorescein-4(5)-carboxylic acid [Khanna & Ullman, (1980) *Anal Biochem.*

20 108:156-161) which had been preactivated with diisopropylcarbodiimide and 1-hydroxybenzotriazole in DMF]. The dimethoxyfluoresceinyl-peptide was then simultaneously deprotected and cleaved from the resin by treatment with trifluoroacetic acid containing 5% each of water and triethylsilane. The dimethoxyfluoresceinyl-peptide was isolated by evaporation, trituration with diethyl ether and filtration. The isolated peptide was reacted with

25 4-(N-maleimido)-fluorescein in DMF containing N,N-diisopropylethylamine, the product purified by RP-HPLC and finally isolated by freeze-drying from aqueous acetic acid. The product was characterised by MALDI-TOF MS and amino acid analysis. The compounds of this invention have been found to be active against TACE (causing at least 50% inhibition) at 50 $\mu$ m and are preferably active at 10 $\mu$ m. In particular, compound 3 caused 50% inhibition at

30 630nm.

### Natural Substrates

The activity of the compounds of the invention as inhibitors of aggrecan degradation may be assayed using methods for example based on the disclosures of E. C. Arner *et al.*, (1998) Osteoarthritis and Cartilage 6:214-228; (1999) Journal of Biological Chemistry, 274 (10), 6594-6601 and the antibodies described therein. The potency of compounds to act as inhibitors against collagenases can be determined as described by T. Cawston and A. Barrett (1979) Anal. Biochem. 99:340-345.

### Inhibition of metalloproteinase activity in cell/tissue based activity

#### Test as an agent to inhibit membrane sheddases such as TNF convertase

10 The ability of the compounds of this invention to inhibit the cellular processing of TNF- $\alpha$  production may be assessed in THP-1 cells using an ELISA to detect released TNF- $\alpha$  essentially as described K. M. Mohler *et al.*, (1994) Nature 370:218-220. In a similar fashion the processing or shedding of other membrane molecules such as those described in N. M. Hooper *et al.*, (1997) Biochem. J. 321:265-279 may be tested using appropriate cell lines and  
15 with suitable antibodies to detect the shed protein.

#### Test as an agent to inhibit cell based invasion

The ability of the compound of this invention to inhibit the migration of cells in an invasion assay may be determined as described in A. Albini *et al.*, (1987) Cancer Research 47:3239-3245.

#### 20 Test as an agent to inhibit whole blood TNF sheddase activity

The ability of the compounds of this invention to inhibit TNF- $\alpha$  production is assessed in a human whole blood assay where LPS is used to stimulate the release of TNF- $\alpha$ . 160 $\mu$ l of heparinized (10Units/ml) human blood obtained from volunteers, was added to the plate and incubated with 20 $\mu$ l of test compound (duplicates), in RPMI1640 + bicarbonate, penicillin,  
25 streptomycin, glutamine and 1% DMSO, for 30 min at 37°C in a humidified (5%CO<sub>2</sub>/95%air) incubator, prior to addition of 20 $\mu$ l LPS (E. coli 0111:B4; final concentration 10 $\mu$ g/ml). Each assay includes controls of neat blood incubated with medium alone or LPS (6 wells/plate of each). The plates are then incubated for 6 hours at 37°C (humidified incubator), centrifuged (2000rpm for 10 min; 4°C), plasma harvested (50-100 $\mu$ l) and stored in 96 well plates at  
30 70°C before subsequent analysis for TNF- $\alpha$  concentration by ELISA.

#### Test as an agent to inhibit in vitro cartilage degradation

The ability of the compounds of this invention to inhibit the degradation of the aggrecan or collagen components of cartilage can be assessed essentially as described by K. M. Bottomley *et al.*, (1997) *Biochem J.* 323:483-488.

#### In vivo assessment

##### 5 Test as an anti-TNF agent

The ability of the compounds of this invention as *in vivo* TNF- $\alpha$  inhibitors is assessed in the rat. Briefly, groups of female Wistar Alderley Park (AP) rats (90-100g) are dosed with compound (5 rats) or drug vehicle (5 rats) by the appropriate route e.g. peroral (p.o.), intraperitoneal (i.p.), subcutaneous (s.c.) 1 hour prior to lipopolysaccharide (LPS) challenge  
10 (30 $\mu$ g/rat i.v.). Sixty minutes following LPS challenge rats are anaesthetised and a terminal blood sample taken via the posterior vena cavae. Blood is allowed to clot at room temperature for 2 hours and serum samples obtained. These are stored at -20°C for TNF- $\alpha$  ELISA and compound concentration analysis.

Data analysis by dedicated software calculates for each compound/dose:

$$15 \text{ Percent inhibition of TNF-}\alpha = \frac{\text{Mean TNF-}\alpha (\text{Vehicle control}) - \text{Mean TNF-}\alpha (\text{Treated})}{\text{Mean TNF-}\alpha (\text{Vehicle control})} \times 100$$

##### Test as an anti-arthritic agent

Activity of a compound as an anti-arthritic is tested in the collagen-induced arthritis (CIA) as defined by D. E. Trentham *et al.*, (1977) *J. Exp. Med.* 146:857. In this model acid  
20 soluble native type II collagen causes polyarthritis in rats when administered in Freund's incomplete adjuvant. Similar conditions can be used to induce arthritis in mice and primates.

##### Pharmaceutical Compositions

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (1), or a pharmaceutically acceptable  
25 salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical  
30 administration as an ointment or cream or for rectal administration as a suppository. The composition may also be in a form suitable for inhalation.

In general the above compositions may be prepared in a conventional manner using



conventional excipients.

The pharmaceutical compositions of this invention will normally be administered to humans so that, for example, a daily dose of 0.5 to 75 mg/kg body weight (and preferably 0.5 to 30 mg/kg body weight) is received. This daily dose may be given in divided doses as  
5 necessary, the precise amount of the compound received and the route of administration depending on the weight, age and sex of the patient being treated and on the particular disease condition being treated according to principles known in the art.

Typically unit dosage forms will contain about 1 mg to 500 mg of a compound of this invention.

10 Therefore in a further aspect of the present invention there is provided a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use in a method of treatment of a warm-blooded animal such as man by therapy. Also provided is a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use in a method of treating a  
15 disease condition mediated by one or more metalloproteinase enzymes and in particular a disease condition mediated by TNF- $\alpha$ . Further provided is a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use in a method of treating inflammatory diseases, autoimmune diseases, allergic/atopic diseases, transplant rejection, graft versus host disease, cardiovascular disease, reperfusion  
20 injury and malignancy in a warm-blooded animal such as man. In particular a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, is provided for use in a method of treating rheumatoid arthritis, Crohn's disease and psoriasis, and especially rheumatoid arthritis in a warm-blooded animal such as man. Also provided is a compound of formula (1), or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, for use in a method of treating a respiratory disorder such as  
25 asthma or COPD in a warm-blooded animal such as man.

According to an additional aspect of the invention there is provided a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use as a medicament. Also provided is a compound of formula (1),  
30 or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use as a medicament in the treatment of a disease condition mediated by one or more metalloproteinase enzymes and in particular a disease condition mediated by TNF- $\alpha$ .

Further provided is a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use as a medicament in the treatment of inflammatory diseases, autoimmune diseases, allergic/atopic diseases, transplant rejection, graft versus host disease, cardiovascular disease, reperfusion injury and malignancy.

- 5 in a warm-blooded animal such as man. In particular a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, is provided for use as a medicament in the treatment of rheumatoid arthritis, Crohn's disease and psoriasis, and especially rheumatoid arthritis in a warm-blooded animal such as man. In addition, a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo*
- 10 hydrolysable ester thereof, is provided for use as a medicament in the treatment of a respiratory disorder such as asthma or COPD in a warm-blooded animal such as man.

- According to this another aspect of the invention there is provided the use of a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment
- 15 of a disease condition mediated by one or more metalloproteinase enzymes and in particular a disease condition mediated by TNF- $\alpha$  in a warm-blooded animal such as man. Also provided is the use of a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of inflammatory diseases, autoimmune diseases, allergic/atopic diseases,
- 20 transplant rejection, graft versus host disease, cardiovascular disease, reperfusion injury and malignancy in a warm-blooded animal such as man. In particular the use of a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, is provided in the manufacture of a medicament in the treatment of rheumatoid arthritis, Crohn's disease and psoriasis, and especially rheumatoid arthritis in a
- 25 warm-blooded animal such as man. In addition, the use of a compound of formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, is provided in the manufacture of a medicament in the treatment of a respiratory disorder such as asthma or COPD in a warm-blooded animal such as man.

- According to a further feature of this aspect of the invention there is provided a
- 30 method of producing a metalloproteinase inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1). According to a further feature of this aspect of the

invention there is provided a method of producing a TACE inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1). According to this further feature of this aspect of the invention there is provided a method of treating autoimmune disease, allergic/atopic diseases, transplant rejection, graft versus host disease, cardiovascular disease, reperfusion injury and malignancy in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1). Also provided is a method of treating rheumatoid arthritis, Crohn's disease and psoriasis, and especially rheumatoid arthritis in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1). Further provided is a method of treating a respiratory disorder such as asthma or COPD in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1).

15 In addition to their use in therapeutic medicine, the compounds of formula (1) and their pharmaceutically acceptable salts are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of cell cycle activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

20 In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

The compounds of this invention may be used in combination with other drugs and therapies used in the treatment of various immunological, inflammatory or malignant disease states which would benefit from the inhibition of TACE.

25 If formulated as a fixed dose such combination products employ the compounds of this invention within the dosage range described herein and the other pharmaceutically-active agent within its approved dosage range. Sequential use is contemplated when a combination formulation is inappropriate.

30

### Examples

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- (i) temperatures are given in degrees Celsius ( $^{\circ}\text{C}$ ); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25 $^{\circ}\text{C}$ ;
- 5 (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30 mm Hg) with a bath temperature of up to 60 $^{\circ}\text{C}$ ;
- (iii) chromatography unless otherwise stated means flash chromatography on silica gel; thin layer chromatography (TLC) was carried out on silica gel plates; where a "Bond Elut" column
- 10 is referred to, this means a column containing 10g or 20g of silica of 40 micron particle size, the silica being contained in a 60ml disposable syringe and supported by a porous disc, obtained from Varian, Harbor City, California, USA under the name "Mega Bond Elut SI". Where an "Isolute<sup>TM</sup> SCX column" is referred to, this means a column containing benzenesulphonic acid (non-endcapped) obtained from International Sorbent Technology Ltd.,
- 15 1st House, Duffryn Industrial Estate, Ystrad Mynach, Hengoed, Mid Glamorgan, UK. Where Flashmaster II is referred to, this means a UV driven automated chromatography unit supplied by Jones;
- (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;
- 20 (v) yields, when given, are for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;
- (vi) when given,  $^1\text{H}$  NMR data is quoted and is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an
- 25 internal standard, determined at 400 MHz using perdeuterio DMSO ( $\text{CD}_3\text{SOCD}_3$ ) as the solvent unless otherwise stated; coupling constants (J) are given in Hz;
- (vii) chemical symbols have their usual meanings; SI units and symbols are used;
- (viii) solvent ratios are given in percentage by volume;
- (ix) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical
- 30 ionisation (APCI) mode using a direct exposure probe; where indicated ionisation was effected by electrospray (ES); where values for m/z are given, generally only ions which

indicate the parent mass are reported, and unless otherwise stated the mass ion quoted is the positive mass ion -  $(M+H)^+$ ;

(x) LCMS (liquid chromatography mass spectrometry) characterisation was performed using a pair of Gilson 306 pumps with Gilson 233 XL sampler and Waters ZMD4000 mass

- 5 spectrometer. The LC comprised water symmetry 4.6x50 column C18 with 5 micron particle size. The eluents were: A, water with 0.05% formic acid and B, acetonitrile with 0.05% formic acid. The eluent gradient went from 95% A to 95% B in 6 minutes. Where indicated ionisation was effected by electrospray (ES); where values for  $m/z$  are given, generally only ions which indicate the parent mass are reported, and unless otherwise stated the mass ion
- 10 quoted is the positive mass ion -  $(M+H)^+$  and


(xi) the following abbreviations are used:

	min	minute(s)
	h	hour(s)
	DIPEA	N,N-diisopropylethylamine
15	DMSO	dimethyl sulphoxide;
	DMF	N-dimethylformamide;
	DCM	dichloromethane;
	NMP	N-methylpyrrolidinone;
	DIAD	diisopropylazodicarboxylate
20	LHMDS or LiHMDS	Lithium bis(trimethylsilyl)amide
	MeOH	methanol
	RT	room temperature
	TFA	trifluoroacetic acid
	EtOH	ethanol
25	EtOAc	ethyl acetate.
	EDTA	ethylenediaminetetraacetic acid
	THF	tetrahydrofuran

#### EXAMPLE 1

R/S-5-[(4-[(2,5-Dimethylbenzyl)oxy]piperidin-1-yl)sulphonyl)methyl]-5-

30 methylimidazolidine-2,4-dione




- To a solution of 4-(2,5-dimethylbenzyloxy)piperidin-1-ylsulphonylpropan-2-one (prepared as described below) (210mg, 0.62mmol) in EtOH (14ml) and water (6ml) was added potassium cyanide (80mg, 1.23mmol) and ammonium carbonate (245mg, 3.10mmol). The mixture was heated at 70°C for 5 h. Additional ammonium carbonate (1g, 12.6mmol) was added and the mixture stirred at RT for 17 h. The mixture was then concentrated to approximately half the volume and extracted with EtOAc (2x10ml). The combined organic layers were partitioned with brine (10ml), dried (MgSO<sub>4</sub>), concentrated and purified by chromatography (10g silica bond elute, eluent 20-100% EtOAc / Hexane) to give R/S-5-[(4-[(2,5-
- 10 dimethylbenzyl)oxy]piperidin-1-yl)sulphonyl)methyl]-5-methylimidazolidine-2,4-dione as a white solid (30mg, 0.07mmol); NMR 1.3 (s, 3H), 1.6 (m, 2H), 1.9 (m, 2H), 2.2 (s, 3H), 2.25 (s, 3H), 3.0 (m, 2H), 3.3 (m, 3H), 3.5 (d, 1H), 3.6 (m, 1H), 4.5 (s, 2H), 7.0 (dd, 1H), 7.05 (dd, 1H), 7.1 (d, 1H), 8.0 (s, 1H), 10.7 (s, 1H); MS (-ve) 408.
- 15 The starting material, 4-(2,5-dimethylbenzyloxy)piperidin-1-ylsulphonylpropan-2-one, was prepared as described below:
- i) To a solution of *tert*-butyl 4-hydroxypiperidin-1-ylcarboxylate (4g, 19.9mmol) in DMF (100ml) at RT was added sodium hydride (796mg, 60% dispersion in oil, 19.9mmol). After 1 h 2,5-dimethylbenzyl chloride (2.94ml, 19.9mmol) was added dropwise. After 16 h
- 20 water was added (5ml) and DMF was removed *in vacuo*. The mixture was partitioned between water (100ml) and DCM (3x200ml) and the combined organic layers were dried (MgSO<sub>4</sub>), concentrated and purified by chromatography (MPLC, eluting with 0→20% EtOAc/DCM) to give *tert*-butyl 4-(2,5-dimethylbenzyloxy)piperidin-1-ylcarboxylate as a green oil (4.15g, 13mmol); NMR 1.4 (m, 11H), 1.8 (m, 2H), 2.2 (d, 6H), 3.0 (m, 2H), 3.6 (m, 3H), 4.4
- 25 (s, 2H), 7.0 (m, 2H), 7.1 (s, 1H); MS: 320.



- ii) To a solution of *tert*-butyl 4-(2,5-dimethylbenzyloxy)piperidin-1-ylcarboxylate (4.1g, 12.85mmol) in DCM (30ml) was added TFA (3ml) and the mixture stirred overnight at RT. TFA (3ml) was added and the mixture stirred at 40°C. After 1 h the mixture was concentrated and the residue azeotroped with toluene to give 4-(2,5-dimethylbenzyloxy)piperidine.TFA salt as a colourless oil (5.52g, 12.85mmol plus a small amount of toluene); NMR 1.7 (m, 2H), 2.0 (m, 2H), 2.2 (s, 3H), 2.25 (s, 3H), 3.0 (m, 2H), 3.2 (m, 2H), 3.65 (m, 1H), 4.45 (s, 2H), 7.0 (m, 2H) and 7.1 (s, 1H); MS: 220.
- iii) To a solution of 4-(2,5-dimethylbenzyloxy)piperidine.TFA salt (5.51g, 12.85mmol plus a small amount of toluene) in DCM (90ml) at 0°C was added triethylamine (8.59ml, 61.6mmol) followed by methanesulphonyl chloride (1.05ml, 13.6mmol) added dropwise over 5 min. The reaction mixture was allowed to warm to RT. After 63 h the mixture was diluted with DCM (90ml), washed with water (50ml), brine (50ml), dried (MgSO<sub>4</sub>) and concentrated to give a light brown oil. The oil was triturated with EtOH (20ml), filtered and washed with cold EtOH and concentrated to give 4-(2,5-dimethylbenzyloxy)piperidinylsulphonylmethane as a white solid (2.63g, 8.0mmol); NMR 1.6 (m, 2H), 1.9 (m, 2H), 2.2 (s, 3H), 2.25 (s, 3H), 2.85 (s, 3H), 3.0 (m, 2H), 3.55 (m, 1H), 4.45 (s, 2H), 7.0 (m, 2H) and 7.1 (s, 1H); MS: 298.
- iv) To a stirred solution of 4-(2,5-dimethylbenzyloxy)piperidin-1-ylsulphonylmethane (500mg, 1.68mmol) in THF (5ml) at 0°C, was added LHMDs (3.6ml, 3.6mmol). After 10 min acetyl chloride (0.14ml, 1.96mmol) was added. After 2 h saturated ammonium chloride (5ml) was added, the reaction warmed to RT and the product extracted with EtOAc (2x10ml). The combined organic layers were partitioned with brine (10ml), dried (MgSO<sub>4</sub>), concentrated and purified by chromatography (10g silica bond elute, eluent 0-50% EtOAc / hexane) to give 4-(2,5-dimethylbenzyloxy)piperidin-1-ylsulphonylpropan-2-one as an oily residue (210mg, 0.62mmol); MS (-ve) 338.

## 25 EXAMPLE 2

R/S-5-[(4-(2-Methylquinolin-4-ylmethoxy)piperidin-1-yl)sulphonyl)methyl]-5-methylimidazolidine-2,4-dione






To a solution of 4-(2-methylquinolin-4-ylmethoxy)piperidin-1-ylsulphonylpropan-2-one (prepared as described below) (206mg, 0.547mmol) in EtOH (5ml) and water (5ml) was added potassium cyanide (69mg, 1.09mmol) and ammonium carbonate (875mg, 3.28mmol). The mixture was heated at 65°C for 2 h. The mixture was then concentrated to approximately  
5 half the volume and extracted with EtOAc (3x10ml). The combined organic layers were partitioned with brine (10ml), dried (MgSO<sub>4</sub>) and concentrated to give a yellow solid. This was recrystallised from hot EtOAc/iso-hexane to give R/S-5-[(4-(2-methylquinolin-4-ylmethoxy)piperidin-1-yl)sulphonyl)methyl]-5-methylimidazolidine-2,4-dione as a white solid (215mg, 0.482mmol); NMR 1.3 (s, 3H), 1.65 (m, 2H), 2.0 (m, 2H), 2.7 (s,  
10 3H), 3.05 (m, 2H), 3.7 (m, 1H), 5.0 (s, 2H), 7.45 (s, 1H), 7.55 (m, 1H), 7.7 (m, 1H), 7.9 (m, 1H), 8.1 (m, 2H); MS (-ve) 445.

The starting material 4-(2-methyl-quinolin-4-yl methoxy)piperidinylsulphonylpropan-2-one, was prepared as described below:

- 15 i) To a stirred suspension of 2-methylquinolin-4-ylcarboxylic acid (4g, 21.4mmol) in THF (100ml) at RT was added lithium aluminium hydride (21.4ml, 1.0M solution in THF, 21.4mmol) dropwise over 20 min. After 16 h water (4ml) was added cautiously followed by 2N NaOH (4ml) and water (12ml). The resulting gelatinous precipitate was filtered off and washed with THF. DCM (200ml) was added to the filtrate and partitioned with saturated  
20 NaHCO<sub>3</sub> (2x75ml). The organic layer was dried (MgSO<sub>4</sub>), concentrated, triturated with DCM and filtered to give 2-methylquinolin-4-ylmethanol as a white powder (858mg, 5mmol). The mother liquours were purified by chromatography (20g silica bond elute, eluent 0→5% EtOH / DCM) to give a further 610mg of product (3.5mmol); NMR 2.6 (s, 3H), 5.0 (d, 2H), 5.5 (t, 1H), 7.4 (s, 1H), 7.5 (t, 1H), 7.7 (t, 1H), 7.9 (m, 2H); MS: 174.
- 25 ii) To a suspension of 2-methylquinolin-4-ylmethanol (100mg, 0.58mmol) in DCM (5ml) at RT was added triethylamine (0.24ml, 1.74mmol). The reaction mixture was then cooled to 0°C and methanesulphonyl chloride (0.05ml, 0.64mmol) was added dropwise. After 10 min the reaction mixture was concentrated, EtOAc (20ml) was added and the organic layer partitioned with brine (10ml), dried (MgSO<sub>4</sub>), concentrated and purified by chromatography  
30 (10g silica bond elute, eluent 5% MeOH / DCM) to give 2-methylquinolin-4-ylmethoxysulphonylmethane (110mg, 0.44mmol); NMR 2.7 (s, 3H), 3.35 (s, 3H), 5.75 (s, 2H), 7.5 (s, 1H), 7.6 (t, 1H), 7.75 (t, 1H), 8.0 (m, 2H); MS: 252.

- iii) To a solution of *tert*-butyl 4-hydroxypiperidin-1-ylcarboxylate (1.75g, 8.73mmol) in DMF (20ml) at 0°C was added sodium hydride (419mg, 60% dispersion in oil, 10.5mmol). After 10 min a solution of 2-methylquinolin-4-ylmethoxysulphonylmethane (2.19g, 8.73mmol) in DMF (10ml) was added dropwise over 5 min at 0°C. After 5 h the mixture was concentrated and the residue taken up in EtOAc (150ml). The organic layer was washed with brine (50ml), dried (Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>), concentrated and purified by chromatography (MPLC, eluting with 75% EtOAc/ hexane) to give *tert*-butyl 4-(2-methylquinolin-4-ylmethoxy)piperidin-1-ylcarboxylate (1.46g, 4.1mmol); MS: 357.
- iv) To a solution of *tert*-butyl 4-(2-methylquinolin-4-ylmethoxy)piperidin-1-ylcarboxylate (1.45g, 4.1mmol) in DCM (10ml) at RT was added TFA (3ml). After 15 h the mixture was concentrated and azeotroped with toluene (x2) to give 4-(2-methylquinolin-4-ylmethoxy)piperidine.di TFA salt (1.97g, 4.1mmol); MS: 257.
- v) To a solution of 4-(2-methylquinolin-4-ylmethoxy)piperidine.di TFA salt (2.49g, 5.2mmol) in DCM (40ml) at 0°C was added triethylamine (4.3ml, 31mmol) followed by methanesulphonyl chloride (0.8ml, 10.3mmol) added dropwise over 1 min. The reaction mixture was allowed to warm to RT. After 15 h the mixture was diluted with DCM (60ml), washed with water (30ml), brine (25ml), concentrated and purified by chromatography (MPLC, eluting with 100% EtOAc) to give 4-(2-methylquinolin-4-ylmethoxy)piperidin-1-ylsulphonylmethane (600mg, 1.8mmol) as a pale yellow solid; NMR 1.6 (m, 2H), 2.0 (m, 2H), 2.65 (s, 3H), 2.85 (s, 3H), 3.0 (m, 2H), 3.3 (m, 2H), 3.7 (m, 1H), 5.0 (s, 2H), 7.4 (s, 1H), 7.5 (t, 1H), 7.7 (t, 1H), 7.9 (d, 1H), 8.0 (d, 1H); MS: 335.
- vi) To a stirred solution of 4-(2-methylquinolin-4-ylmethoxy)piperidin-1-ylsulphonylmethane (400mg, 1.06mmol) in dry THF (10ml) at approximately -16°C, was added LHMDS (2.63ml, 2.34mmol). After 30 min EtOAc (0.1ml, 1.06mmol) was added and the reaction warmed to RT. After 2 h saturated ammonium chloride (10ml) was added, the reaction warmed to RT and the product extracted with EtOAc (3x20ml). The combined organic layers were partitioned with brine (10ml), dried (MgSO<sub>4</sub>) and concentrated to give the crude product (0.36g) as a yellow oil. This was titrated with iso-hexane to give 4-(2-methylquinolin-4-yl methoxy)piperidin-1-ylsulphonylpropan-2-one as a white solid (206mg, 0.547mmol); NMR 1.65 (m, 2H), 1.9 (m, 2H), 2.2 (s, 2H), 2.6 (s, 3H), 3.0 (m, 2H), 3.35 (m, 2H), 3.6 (m, 1H), 4.9 (s, 2H), 7.4 (s, 1H), 7.5 (t, 1H), 7.6 (t, 1H), 7.85 (d, 1H), 8.0 (d, 1H); MS (+ve) 377.

**EXAMPLE 3****5-[2-({4-[(2-Methylquinolin-4-yl)methoxy]piperidin-1-yl}sulphonyl)ethyl]imidazolidine-2,4-dione**

- 5 To a stirred solution of 2-methyl-4-[(piperidin-4-yloxy)methyl]quinoline. di TFA salt (example 2 step iv)) (200mg, 0.78mmol) in DCM (20ml) under argon, was added DIPEA (0.4ml, 2.35mmol) followed by 2-(2,5-dioxo-4-imidazolidinyl)-1-ethanesulphonyl chloride (see below) (305mg, 1.56mmol). This was stirred for 2 h. Water (ca. 20ml) was added and the resultant emulsion was filtered through celite. The organic phase was dried (MgSO<sub>4</sub>) and
- 10 evaporated, and the residue was purified via chromatography (silica, 1-5% MeOH/EtOAc) to give 5-[2-({4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl}sulphonyl)ethyl]imidazolidine-2,4-dione(175mg, 0.39mmol) as a white solid; NMR 1.7 (m, 2H), 1.9 (m, 3H), 2.1 (m, 1H), 2.7 (s, 3H), 3.2 (m, 4H), 3.5 (m, 2H), 3.8 (m, 3H), 4.2 (m, 1H), 5.05 (s, 2H), 7.4 (s, 1H), 7.5 (t, 1H), 7.6 (s, 1H), 7.7 (t, 1H), 7.9 (d, 1H), 8.1 (d, 1H); MS 447 (MH<sup>+</sup>).

15


The starting material 2-(2,5-dioxo-4-imidazolidinyl)-1-ethanesulphonyl chloride was prepared as follows:

- i) Commercially available RS homocysteine (0.18mol) was suspended in water (25ml). Potassium cyanate (1.5g, 0.2mol) was added and the mixture was stirred at 100°C for 45 min.
- 20 After partial cooling, 10% HCl (10ml) was added and the mixture stirred at 100°C for 50 min. The mixture was placed in the fridge overnight, and the resultant crystals were filtered, washed successively with water and dried *in vacuo* to afford 5-(2-{[2-(2,5-dioxo-4-imidazolidinyl)ethyl]disulphanyl}ethyl)-2,4-imidazolidinedione; LCMS (APCI) m/z 319.1 (MH<sup>+</sup>).
- 25 ii) Through a suspension of 5-(2-{[2-(2,5-dioxo-4-imidazolidinyl)ethyl]disulphanyl}ethyl)-2,4-imidazolidinedione (6.9 mol) in a mixture of AcOH (25ml) and water (2ml) stirred vigorously at 0°C, was bubbled chlorine gas for 15 min

(until all precipitate dissolved) at a maximum temperature of 5°C. The mixture was further stirred for 15 min, evaporated to a small volume *in vacuo* (maximum temperature 30°C), dissolved in DCM (50ml), shaken carefully with saturated NaHCO<sub>3</sub> (ca 25 ml) and then 10% sodium thiosulphate, dried, evaporated and crystallised from THF-hexane (Lora-Tamayo, M. 5 *et al.*, 1968, *An. Quim.*, 64(6):591-606) to afford 2-(2,5-dioxo-4-imidazolidinyl)-1-ethanesulphonyl chloride; NMR 2.55 (m, 1.1H), 2.65 (m, 1.8H), 2.70 (m, 1H), 4.55 (m, 1H).

#### EXAMPLE 4

5-[2-[(4-[(2-Methylquinolin-4-yl)oxy]methyl)piperidin-1-yl)sulphonyl]ethyl]imidazolidine-2,4-dione



10

To a solution of 2-methyl-4-(piperidin-4-ylmethoxy)quinoline (hydrochloric acid salt) (50mg, 0.17mmol) in DMF (5ml) was added DIPEA (0.2ml, 1.02mmol) followed by 2-(2,5-dioxo-4-imidazolidinyl)-1-ethanesulphonyl chloride (see example 3) (76mg, 0.34mmol). The reaction mixture was stirred for 3 h and then partitioned between EtOAc and water. The organic phase 15 was dried and evaporated to give 5-[2-[(4-[(2-methylquinolin-4-yl)oxy]methyl)piperidin-1-yl)sulphonyl]ethyl]imidazolidine-2,4-dione (25mg, 0.056mmol); NMR 1.4 (m, 2H), 1.9 (m, 3H), 2.15 (m, 2H), 2.6 (s, 3H), 3.2 (m, 2H), 3.7 (m, 2H), 4.2 (m, 3H), 6.9 (s, 1H), 7.45 (m, 1H), 7.7 (m, 1H), 7.8 (d, 1H), 8.1 (d, 1H), 10.4 (s, 1H); MS 447.

20 The starting material 2-methyl-4-(piperidin-4-ylmethoxy)quinoline (hydrochloric acid salt) was prepared as follows:


i) *Tert*-butyl 4-(hydroxymethyl)piperidin-1-ylcarboxylate (3.0g) was dissolved in DMF (30ml) with stirring. Sodium hydride (60% in mineral oil, 558mg) was then added and the mixture stirred at 80°C, under argon, for 30 min. A solution of 4-chloroquinaldine (2.5g) in 25 DMF (20ml) was added, followed by potassium fluoride (100mg) and the mixture stirred at 80°C for 5 h. The mixture was concentrated *in vacuo* and the residue partitioned between EtOAc (100ml) and water (100ml). The aqueous phase was extracted with EtOAc (100ml)

and the combined organic layers were washed with brine (100ml), dried ( $\text{MgSO}_4$ ), concentrated *in vacuo* and purified on a bondelut cartridge, eluting with a gradient of 10-75% EtOAc/iso-hexane. The compound obtained was purified on a second identical cartridge, eluting with a gradient of 0-40% EtOAc/iso-hexane to give *tert*-butyl 4-[[[(2-methylquinolin-4-yl)oxy]methyl]piperidin-1-yl]carboxylate as a white solid (2.27g); NMR 1.25 (m, 2H), 1.40 (s, 9H), 1.82 (m, 2H), 2.10 (m, 1H), 2.55 (s, 3H), 2.75 (m, 2H), 4.00 (m, 2H), 4.10 (d, 2H), 6.92 (s, 2H), 7.45 (m, 1H), 7.65 (t, 1H), 7.82 (d, 1H), 8.05 (d, 1H); MS 357 ( $\text{MH}^+$ ).

ii) *Tert*-butyl 4-[[[(2-methylquinolin-4-yl)oxy]methyl]piperidin-1-yl]carboxylate (2.27g) was stirred in a mixture of concentrated hydrochloric acid (12.5ml) and 1,4-dioxane (25ml) for 16 h. The mixture was then concentrated *in vacuo*, azeotroped with toluene (3x30ml) and dried *in vacuo* to give 2-methyl-4-(piperidin-4-ylmethoxy)quinoline (hydrochloric acid salt) as an off-white solid (2.04g); NMR 1.75 (m, 2H), 2.00 (m, 2H), 2.30 (m, 1H), 2.90 (s, 3H), 2.95 (m, 2H), 3.35 (m, 2H), 4.40 (m, 2H), 7.50 (s, 1H), 7.80 (t, 1H), 8.10 (t, 1H), 8.30 (d, 1H), 8.40 (d, 1H), 9.15 (m, 2H); MS 257 ( $\text{MH}^+$ ).

#### 15 EXAMPLE 5

**5-Methyl-5-[[[(4-[[[(2-methylquinolin-4-yl)oxy]methyl]piperidin-1-yl)sulphonyl]methyl]imidazolidine-2,4-dione**



An analogous method to that described in example 4 was used except that 2-(2,5-dioxo-4-imidazolidinyl)-1-ethanesulphonyl chloride was replaced with [4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride. 5-methyl-5-[[[(4-[[[(2-methylquinolin-4-yl)oxy]methyl]piperidin-1-yl)sulphonyl]methyl]imidazolidine-2,4-dione was obtained as a white solid (53mg); NMR 1.32 (s, 3H), 1.45 (m, 2H), 1.95 (m, 2H), 2.12 (s, 1H), 2.85 (m, 5H), 3.35 (m, 1H), 3.50 (m, 1H), 3.60 (t, 2H), 4.40 (d, 2H), 7.52 (s, 1H), 7.83 (m, 1H), 8.00 (s, 1H), 8.08 (m, 2H), 8.35 (m, 1H), 10.70 (s, 1H); MS 447 ( $\text{MH}^+$ ).


The starting material [4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride was prepared as follows:

- i) To a steel vessel charged with EtOH (315ml) and water (135ml) was added benzylthioacetone (31.7g, 0.175 mol), potassium cyanide (22.9g, 0.351 mol) and ammonium carbonate (84.5g, 0.879 mol). The closed reaction vessel was kept at 90 °C under vigorous stirring for 3 h. The reaction vessel was then cooled with ice-water (30 min), the resultant yellowish slurry evaporated to dryness and the solid residue partitioned between water (400ml) and EtOAc (700ml) and separated. The aqueous phase was extracted with EtOAc (300ml). The combined organic phases were washed with saturated brine (150ml), dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to dryness. (Crystallisation was assisted by the addition of DCM (300ml) to the oil). Evaporation gave 5-methyl-5-  
10 {[(phenylmethyl)thio]methyl}imidazolidine-2,4-dione as a slightly yellowish powder (43.8g, 90%); NMR 1.29 (3H, s), 3.76 (2H, s), 2.72, 2.62 (1H each, ABq,  $J=14.0$  Hz), 7.35-7.20 (5H, m), 8.00 (1H, s), 10.74 (1H, s); (MH<sup>+</sup>)  $m/z$  251.1.
- ii) 5-Methyl-5-[(phenylmethyl)thio]methyl}imidazolidine-2,4-dione (42.6g; 0.17mol) was dissolved in a mixture of AcOH (450ml) and water (50ml). The mixture was immersed in an ice/water bath and chlorine gas was bubbled through the solution, such that the temperature was maintained below 15 °C. After 25 min the solution became yellow-green in colour and a sample was withdrawn for LCMS and HPLC analysis. It showed that the starting material had been consumed. The yellow clear solution was stirred for 30 min and an opaque solution /slurry was formed. The solvent was removed *in vacuo* at 37°C, the resultant yellowish solid  
20 was suspended in toluene (400ml) and solvent removed *in vacuo*. This was repeated. The crude product was then suspended in iso-hexane (400ml) and warmed to 40°C while stirring. The slurry was then allowed to cool to RT before the insoluble product was removed by filtration, washed with iso-hexane (6x100ml) and dried under reduced pressure at 50°C overnight. This gave [4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride as a  
25 slightly yellow powder (36.9g, 95%); Purity by HPLC = 99%, NMR supported purity; NMR (THF- $d_8$ ) 9.91 (1H, bs), 7.57 (1H, s), 4.53, 4.44 (1H each, ABq,  $J=14.6$ Hz), 1.52 (s, 3H,  $\text{CH}_3$ );  $^{13}\text{C}$  NMR (THF- $d_8$ )  $\delta$  174.96, 155.86, 70.96, 61.04, 23.66.

#### EXAMPLE 6

5-Ethyl-5-[(4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)methyl]imidazolidine-2,4-dione  
30






2-Methyl-4-[(piperidin-4-yloxy)methyl]quinoline di TFA salt (example 2 step iv)) (100mg, 0.39mmol) was dissolved in DMF (5ml) under argon. DIPEA (0.2ml, 1.18mmol) was added, followed by [4-ethyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride (145mg, 5 0.59mmol) and the mixture was stirred for 20 h. The reaction mixture was then partitioned between EtOAc and water, the organic phases separated, washed with water, dried and evaporated, and the residue triturated with ether to give 5-ethyl-5-[(4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)methyl]imidazolidine-2,4-dione as a white solid (10mg, 0.02mmol); NMR 0.9 (m, 3H), 1.7 (m, 4H), 2.0 (m, 2H), 2.7 (s, 3H), 3.1 (m, 2H), 3.3 (m, 3H), 10 3.8 (m, 1H), 5.1 (s, 2H), 7.4-7.6 (m, 3H), 7.7 (m, 1H), 7.9 (d, 1H), 8.0 (d, 1H).

The starting material [4-ethyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride was prepared using an analogous method to that used in example 5 to prepare [4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride except that benzylthioacetone was 15 replaced with 1-(benzylthio)butan-2-one (Tetrahedron Letters (1998), 39(20), 3189-3192.); NMR (THF-d8) 0.9 (3H, t), 1.9 (2H, m), 4.4 (1H, d), 4.5 (1H, d), 7.4 (1H, s), 9.9 (1H, s).

#### EXAMPLE 7

**5-Methyl-5-[2-((4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)ethyl]imidazolidine-2,4-dione**



20

An analogous method to that described in example 6 was used except that [4-ethyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride was replaced with [4-methyl-2,5-dioxoimidazolidin-4-yl]ethanesulphonyl chloride to afford 5-methyl-5-[2-((4-[(2-




methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)ethyl]imidazolidine-2,4-dione as an off white solid; NMR 1.29 (s, 3H), 1.70 (m, 2H), 1.97 (m, 4H), 2.89 (m, 4H), 3.08 (m, 3H), 3.44 (m, 2H), 3.78 (m, 1H), 5.23 (s, 2H), 7.81 (m, 2H), 8.01 (m, 2H), 8.16 (d, 1H), 8.37 (d, 1H), 10.73 (s, 1H); MS 461 (MH<sup>+</sup>).

5

The starting material [4-methyl-2,5-dioxoimidazolidin-4-yl]ethanesulphonyl chloride was prepared by an analogous method to that described in example 5 to prepare [4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride except that benzylthioacetone was replaced with 1-(benzylthio)butan-3-one (Angewandte Chemie, International Edition (2000), 10 39(23), 4316-4319); NMR (THF-d<sub>8</sub>) 1.4 (s, 3H), 2.25 (m, 1H), 2.35 (m, 1H), 3.85 (m, 1H), 4.0 (m, 1H), 7.1 (s, 1H), 9.8 (s, 1H).

### EXAMPLE 8

**5-Ethyl-5-[2-({4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)ethyl]imidazolidine-2,4-dione**



15


An analogous method to that described in example 6 was used except that [4-ethyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride was replaced with [4-ethyl-2,5-dioxoimidazolidin-4-yl]ethanesulphonyl chloride to afford 5-ethyl-5-[2-({4-[(2-methylquinolin-4-yl)methoxy]piperidin-1-yl)sulphonyl)ethyl]imidazolidine-2,4-dione as an off white solid; NMR 0.77 (t, 3H), 1.57-1.77 (m, 4H), 1.88-2.06 (m, 4H), 2.86 (m, 4H), 3.10 (m, 3H), 3.44 (m, 2H), 3.77 (m, 1H), 5.21 (s, 2H), 7.79 (m, 2H), 7.90 (s, 1H), 8.00 (t, 1H), 8.12 (d, 1H), 8.23 (d, 1H), 10.73 (s, 1H), MS 475 (MH<sup>+</sup>).

The starting material [4-ethyl-2,5-dioxoimidazolidin-4-yl]ethanesulphonyl chloride was prepared by an analogous method to that described in example 5 to prepare [4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride except that benzylthioacetone was replaced with 1-(benzylthio)pentan-3-one (Chemical & Pharmaceutical Bulletin (1997), 25

45(5), 778-785.); NMR (THF-d<sub>8</sub>) 0.9 (t, 3H), 1.7 (m, 1H), 1.9 (m, 1H), 2.2 (m, 1H), 2.35 (m, 1H), 3.9 (m, 1H), 4.0 (m, 1H), 7.1 (s, 1H), 9.8 (s, 1H).

### EXAMPLE 9

(5S)-5-Methyl-5-{4-[(2-methylquinolin-4-yl)methoxymethyl]piperidylsulphonylmethyl}-2,4-dioxoimidazolidine



An analogous method to that described in example 6 was used except that 2-methyl-4-[(piperidin-4-yloxy)methyl]quinoline.diTFA salt was replaced with 4-[(2-methylquinolin-4-yl)methoxymethyl]piperidine and [4-ethyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride replaced with (4S)-(4-methyl-2,5-dioxoimidazolidin-4-yl)methanesulphonyl chloride to afford (5S)-5-methyl-5-{4-[(2-methylquinolin-4-yl)methoxymethyl]piperidylsulphonylmethyl}-2,4-dioxoimidazolidine as an off white solid; NMR 1.1 (m, 2H), 1.2 (m, 3H), 1.7-1.8 (m, 3H), 2.6 (s, 3H), 2.7-2.8 (m, 2H), 3.3 (m, partly obscured by H<sub>2</sub>O, 2H), 3.3-3.5 (m, 4H), 4.9 (s, 2H), 7.4 (s, 1H), 7.5 (t, 1H), 7.7 (t, 1H), 7.9-8.1 (m, 3H), 10.7 (s, 1H); MS 461 (MH<sup>+</sup>).

The starting material 4-[(2-methylquinolin-4-yl)methoxymethyl]piperidine was prepared as follows :-

i) To a stirred solution of 2-methyl-4-hydroxymethylquinoline (2.22g) in DMF (40ml) was added a 60% suspension of sodium hydride in mineral oil (620mg). After 15 min, *tert*-butyl 4-[[[(4-methylphenyl)sulphonyl]oxy]methyl]piperidin-1-ylcarboxylate (4.7g) (Preparation of quinazolinyl ureas, thioureas and guanidines for use in the prevention or treatment of T cell mediated diseases or medical conditions; Crawley, McKerrecher, Poyser, Hennequin and Lambert (Astrazeneca UK Limited, UK; Zeneca Pharma S.A.) WO 0104102 169 pp) was added and the mixture stirred at 20°C for 18 h. The mixture was quenched carefully with water (100ml) and extracted repeatedly with EtOAc. The combined EtOAc extracts were washed with water, brine, dried and evaporated to an oil. This was chromatographed on silica in EtOAc-isohexane mixtures affording *tert*-butyl 4-[[[(2-methylquinolin-4-yl)methoxy]methyl] piperidin-1-ylcarboxylate

(1.2g) as an oil; NMR 1.1-1.3 (m, 2H), 1.35 (s, 9H), 1.7-1.9 (m, 3H), 2.6-2.8 (s, m, 5H), 3.45 (d, 2H), 4.0-4.2 (m, 2H), 4.9 (s, 2H), 7.3 (s, 1H), 7.45 (t, 1H), 7.65 (t, 1H), 7.8 (d, 1H), 8.05 (d, 1H); MS 371.2 (MH<sup>+</sup>).

ii) 4M Dioxan-HCl (40ml) was added to a solution of *tert*-butyl 4-[[2-methylquinolin-4-yl)methoxy]methyl]piperidin-1-ylcarboxylate (1.0g) in dioxan (5ml). MeOH (4ml) was added and the mixture stirred at 20°C for 2 h. The solvents were evaporated, the residue dissolved in DCM (20ml), washed with water, NaHCO<sub>3</sub> and brine, dried and evaporated to a gum (1.0g); NMR 1.2-1.3 (m, 2H), 1.7-1.9 (m, 3H), 2.5-2.75 (m, 6H), 3.15 (d, 2H), 3.45 (d, 2H), 4.9 (s, 2H), 7.3 (s, 1H), 7.45 (t, 1H), 7.65 (t, 1H), 7.9 (d, 1H), 8.05 (d, 1H).

10

The starting material (4S)-(4-methyl-2,5-dioxoimidazolidin-4-yl)methanesulphonyl chloride was prepared as follows :-

i) A steel vessel was charged with EtOH (315ml) and water (135ml), and benzylthioacetone (31.7g, 0.175mol), potassium cyanide (22.9g, 0.351mol) and ammonium carbonate (84.5g, 0.879mol) were added. The closed reaction vessel was heated to 90 °C and stirred vigorously for 3h. The reaction vessel was cooled with ice-water for 30 min, the yellowish slurry evaporated to dryness, the solid residue partitioned between water (400ml) and EtOAc (700ml) and then separated. The aqueous phase was extracted with EtOAc (300ml) and the combined organic phases were washed with saturated brine (150ml), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to dryness. Crystallisation was assisted by the addition of DCM (300ml). Evaporation gave 5-methyl-5-[[phenylmethyl]thio]methyl]imidazolidine-2,4-dione as a slightly yellowish powder (43.8g, 90%); LC-MS (APCI) m/z 251.1 (MH<sup>+</sup>); NMR 10.74 (1H, s), 8.00 (1H, s), 7.35-7.20 (5H, m), 3.76 (2H, s), 2.72, 2.62 (1H each, ABq, *J*=14.0 Hz), 1.29 (3H, s); <sup>13</sup>C NMR DMSO-d<sub>6</sub> 177.30, 156.38, 138.11, 128.74, 128.24, 126.77, 62.93, 37.96, 36.39, 23.15.

ii) (5S)-5-Methyl-5-[[phenylmethyl]thio]methyl]imidazolidine-2,4-dione was prepared by chiral separation of the racemic material using a 250mm x 50mm column on a Dynamic Axial Compression Preparative HPLC system. The stationary phase used was CHIRALPAK AD and other parameters were: MeOH (eluent), 89ml/min (flow), RT, 220nm (UV), 150mg/ml (sample concentration) and 20ml (injection volume). The retention time for this enantiomer under these conditions was 6 min. Analysis of chiral purity was made using a 250mm x 4.6mm CHIRALPAK-AD column from Daicel with parameters: 0.5ml/min (flow),

EtOH (eluent), 220nm (UV), RT. The retention time for this enantiomer under these conditions was 9.27 min. Chiral purity was estimated to >99% ee.

LC-MS (APCI)  $m/z$  251.1 (MH<sup>+</sup>);  $[\alpha]_D = -30.3^\circ$  ( $c=0.01\text{g/ml}$ , MeOH,  $T=20^\circ\text{C}$ ); NMR 10.74 (1H, s), 8.00 (1H, s), 7.35-7.20 (5H, m), 3.76 (2H, s), 2.72, 2.62 (1H each, ABq,  $J=14.0\text{ Hz}$ ),  
5 1.29 (3H, s);  $^{13}\text{C}$  NMR (DMSO- $d_6$ ) 177.30, 156.28, 138.11, 128.74, 128.24, 126.77, 62.93, 37.96, 36.39, 23.15.

(5*R*)-5-methyl-5-[[*(phenylmethyl)*thio]methyl]imidazolidine-2,4-dione was similarly prepared by chiral separation of the racemic material using a 250mm x 50mm column on a Dynamic Axial Compression Preparative HPLC system. The stationary phase used was

- 10 CHIRALPAK AD and other parameters were: MeOH (eluent), 89ml/min (flow), RT, 220nm (UV), 150mg/ml (sample concentration), 20ml (injection volume). The retention time for this enantiomer under these conditions was 10 min. Analysis of chiral purity was made using a 250mm x 4.6mm CHIRALPAK-AD column from Daicel with parameters: 0.5ml/min (flow), EtOH (eluent), 220nm (UV), RT. The retention time for this enantiomer under these  
15 conditions was 17.81 min. Chiral purity was estimated to >99% ee.

LC-MS (APCI)  $m/z$  251.0 (MH<sup>+</sup>);  $[\alpha]_D = +30.3^\circ$  ( $c=0.01\text{g/ml}$ , MeOH,  $T=20^\circ\text{C}$ ); NMR 10.74 (1H, s), 8.00 (1H, s), 7.35-7.20 (5H, m), 3.76 (2H, s), 2.72, 2.62 (1H each, ABq,  $J=14.0\text{ Hz}$ ), 1.29 (3H, s);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$ : 177.31, 156.30, 138.11, 128.74, 128.25, 126.77, 62.94, 37.97, 36.40, 23.16.


- 20 iii) (5*S*)-5-Methyl-5-[[*(phenylmethyl)*thio]methyl]imidazolidine-2,4-dione (42.6g, 0.17mol) was dissolved in a mixture of AcOH (450 ml) and water (50ml). The mixture was immersed in an ice/water bath and chlorine gas was bubbled through the solution, the flow of gas being adjusted to maintain the temperature at below  $15^\circ\text{C}$ . After 25 min the solution became yellow-green in colour and a sample was withdrawn for LC/MS and HPLC analysis.  
25 It showed that starting material had been consumed. The resultant clear yellow solution was stirred for 30 min and an opaque solution /slurry was formed. The solvent was removed on a rotary evaporator at  $37^\circ\text{C}$ , the resultant yellowish solid suspended in toluene (400ml) and solvent removed on the same rotary evaporator. This was repeated once more. The crude product was then suspended in iso-hexane (400ml) and warmed to  $40^\circ\text{C}$  while stirring. The  
30 slurry was allowed to cool to RT before the insoluble product was removed by filtration, washed with iso-hexane (6x100ml), and dried under reduced pressure at  $50^\circ\text{C}$  over night. This gave [(4*S*)-4-methyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride as a slightly

yellow powder (36.9g, 95%); purity by HPLC = 99%, NMR supported that purity;  $[\alpha]_D = -12.4^\circ$  ( $c=0.01\text{g/ml}$ , THF,  $T=20^\circ\text{C}$ ); NMR (THF- $d_8$ ) 9.91 (1H, bs), 7.57 (1H, s), 4.53, 4.44 (1H each, ABq,  $J=14.6\text{Hz}$ ), 1.52 (s, 3H,  $\text{CH}_3$ );  $^{13}\text{C}$  NMR (THF- $d_8$ ) 174.96, 155.86, 70.96, 61.04, 23.66.

- 5 [(4*R*)-4-Methyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride can be prepared by the same method;  $[\alpha]_D = +12.8^\circ$  ( $c=0.01\text{g/ml}$ , THF,  $T=20^\circ\text{C}$ ), NMR (THF- $d_8$ ) 9.91 (1H, brs), 7.57 (1H, s), 4.53, 4.44 (1H each, ABq,  $J=14.6\text{Hz}$ ), 1.52 (s, 3H,  $\text{CH}_3$ );  $^{13}\text{C}$  NMR (THF- $d_8$ ) 174.96, 155.84, 70.97, 61.04, 23.66.

#### EXAMPLE 10

- 10 (5*S*)-5-Ethyl-5-{4-[(2-methylquinolin-4-yl)methoxymethyl]piperidylsulphonylmethyl}-2,4-dioxoimidazolidine



- An analogous method to that described in example 6 was used except that 2-methyl-4-[(piperidin-4-yloxy)methyl]quinoline was replaced with 4-[(2-methylquinolin-4-yl)methoxymethyl]piperidine and [4-ethyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride was replaced with (4*S*)-2-(4-ethyl-2,5-dioxoimidazolidin-4-yl)methanesulphonyl chloride to afford (5*S*)-5-ethyl-5-{4-[(2-methylquinolin-4-yl)methoxymethyl]piperidylsulphonylmethyl}-2,4-dioxoimidazolidine as an off white solid; NMR 0.75 (t, 3H), 1.1-1.3 (m, 2H), 1.6 (q, 2H), 1.7-1.85 (m, 3H), 2.65 (s, 3H), 2.7-2.9 (m, 2H), 3.2-3.3 (m, 2H), 3.3-3.6 (m, 4H), 4.9 (s, 2H), 7.4 (s, 1H), 7.5 (t, 1H), 7.7 (t, 1H), 7.8 (m, 2H), 8.0 (d, 1H), 10.7 (s, 1H); MS 473 (MH $^-$ ).

- The starting material 2-methyl-4-[(piperidin-4-yloxy)methyl]quinoline was prepared as described in example 9 and (4*S*)-2-(4-ethyl-2,5-dioxoimidazolidin-4-yl)methanesulphonyl chloride was prepared as described in example 6. Separation of the isomers was achieved as described in example 9.


(5*S*)-5-Ethyl-5-[[[(phenylmethyl)thio]methyl]imidazolidine-2,4-dione: the retention time for this enantiomer was 4.1 min; NMR 0.77 (t, 3H), 1.59 (m, 2H), 2.67 (q, 2H), 3.76 (s, 2H), 7.24 (m, 1H), 7.31 (m, 4H), 7.89 (s, 1H), 10.72 (s, 1H).

[(4*S*)-4-Ethyl-2,5-dioxoimidazolidin-4-yl]methanesulphonyl chloride; NMR (THF- $d_8$ ) 0.96 (s, 3H), 1.90 (m, 2H); 4.49 (d, 1H), 4.59 (d, 1H), 7.54 (s, 1H), 9.97 (s, 1H).

**CLAIMS**

We claim:

1. A compound of formula (1) or a pharmaceutically acceptable salt thereof:



formula (1)

wherein:

 $Y^1$  and  $Y^2$  are independently O or S;z is  $NR^8$ , O or S;

10 n is 0 or 1;

W is  $NR^1$ ,  $CR^1R^2$  or a bond;

m is 0 or 1;

D is hydrogen,  $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl or fluoro;X is  $-(CR^{12}R^{13})_t-Q-(CR^{14}R^{15})_u-$  where t and u are independently 0 or 1 and Q is O, S, SO or15  $SO_2$ ;

B is a group selected from aryl, heteroaryl and heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl,

trifluoromethoxy, halo, cyano,  $C_{1-4}$ alkyl (optionally substituted by  $R^9$  or  $C_{1-4}$ alkoxy or one or more halo),  $C_{2-4}$ alkenyl (optionally substituted by halo or  $R^9$ ),  $C_{2-4}$ alkynyl (optionally20 substituted by halo or  $R^9$ ),  $C_{3-6}$ cycloalkyl (optionally substituted by  $R^9$  or one or more halo), $C_{5-6}$ cycloalkenyl (optionally substituted by halo or  $R^9$ ), aryl (optionally substituted by halo or $C_{1-4}$ alkyl), heteroaryl (optionally substituted by halo or  $C_{1-4}$ alkyl), heterocyclyl (optionallysubstituted by  $C_{1-4}$ alkyl),  $-SR^{11}$ ,  $-SOR^{11}$ ,  $-SO_2R^{11}$ ,  $-SO_2NR^9R^{10}$ ,  $-NR^9SO_2R^{11}$ , - $NHCONR^9R^{10}$ ,  $-OR^9$ ,  $-NR^9R^{10}$ ,  $-CONR^9R^{10}$  and  $-NR^9COR^{10}$ ; or B is  $C_{2-4}$ alkenyl or  $C_{2-}$ 25  $C_{4}$ alkynyl, each being optionally substituted by a group selected from  $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl,

aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more halo,



- nitro, cyano, trifluoromethyl, trifluoromethoxy,  $-\text{CONHR}^9$ ,  $-\text{CONR}^9\text{R}^{10}$ ,  $-\text{SO}_2\text{R}^{11}$ ,  $-\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $-\text{NR}^9\text{SO}_2\text{R}^{11}$ ,  $\text{C}_{1-4}\text{alkyl}$  or  $\text{C}_{1-4}\text{alkoxy}$ ; with the provisos that:
- when  $n$  is 1 and  $W$  is  $\text{NR}^1$ ,  $\text{CR}^1\text{R}^2$  or a bond; or when  $n$  is 0 and  $W$  is  $\text{CR}^1\text{R}^2$ ; then  $B$  is a group selected from aryl, heteroaryl and heterocyclyl, where each group is optionally
- 5 substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo, cyano,  $\text{C}_{1-4}\text{alkyl}$  (optionally substituted by  $\text{R}^9$  or  $\text{C}_{1-4}\text{alkoxy}$  or one or more halo),  $\text{C}_{2-4}\text{alkenyl}$  (optionally substituted by halo or  $\text{R}^9$ ),  $\text{C}_{2-4}\text{alkynyl}$  (optionally substituted by halo or  $\text{R}^9$ ),  $\text{C}_{3-6}\text{cycloalkyl}$  (optionally substituted by  $\text{R}^9$  or one or more halo),  $\text{C}_{5-6}\text{cycloalkenyl}$  (optionally substituted by halo or  $\text{R}^9$ ), aryl (optionally substituted by halo or
- 10  $\text{C}_{1-4}\text{alkyl}$ ), heteroaryl (optionally substituted by halo or  $\text{C}_{1-4}\text{alkyl}$ ), heterocyclyl (optionally substituted by  $\text{C}_{1-4}\text{alkyl}$ ),  $-\text{SR}^{11}$ ,  $-\text{SOR}^{11}$ ,  $-\text{SO}_2\text{R}^{11}$ ,  $-\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $-\text{NR}^9\text{SO}_2\text{R}^{11}$ ,  $-\text{NHCONR}^9\text{R}^{10}$ ,  $-\text{OR}^9$ ,  $-\text{NR}^9\text{R}^{10}$ ,  $-\text{CONR}^9\text{R}^{10}$  and  $-\text{NR}^9\text{COR}^{10}$ ; or  $B$  is  $\text{C}_{2-4}\text{alkenyl}$  or  $\text{C}_{2-4}\text{alkynyl}$ , each being optionally substituted by a group selected from  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{3-6}\text{cycloalkyl}$ , aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more halo,
- 15 nitro, cyano, trifluoromethyl, trifluoromethoxy,  $-\text{CONHR}^9$ ,  $-\text{CONR}^9\text{R}^{10}$ ,  $-\text{SO}_2\text{R}^{11}$ ,  $-\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $-\text{NR}^9\text{SO}_2\text{R}^{11}$ ,  $\text{C}_{1-4}\text{alkyl}$  or  $\text{C}_{1-4}\text{alkoxy}$ ; and
- when  $n$  is 0 and  $W$  is  $\text{NR}^1$  or a bond; then  $B$  is a group selected from bicyclic aryl, bicyclic heteroaryl and bicyclic heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo,
- 20 cyano,  $\text{C}_{1-4}\text{alkyl}$  (optionally substituted by  $\text{R}^9$  or  $\text{C}_{1-4}\text{alkoxy}$  or one or more halo),  $\text{C}_{2-4}\text{alkenyl}$  (optionally substituted by halo or  $\text{R}^9$ ),  $\text{C}_{2-4}\text{alkynyl}$  (optionally substituted by halo or  $\text{R}^9$ ),  $\text{C}_{3-6}\text{cycloalkyl}$  (optionally substituted by  $\text{R}^9$  or one or more halo),  $\text{C}_{5-6}\text{cycloalkenyl}$  (optionally substituted by halo or  $\text{R}^9$ ), aryl (optionally substituted by halo or  $\text{C}_{1-4}\text{alkyl}$ ), heteroaryl (optionally substituted by halo or  $\text{C}_{1-4}\text{alkyl}$ ), heterocyclyl (optionally substituted by  $\text{C}_{1-4}\text{alkyl}$ ),
- 25  $-\text{SR}^{11}$ ,  $-\text{SOR}^{11}$ ,  $-\text{SO}_2\text{R}^{11}$ ,  $-\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $-\text{NR}^9\text{SO}_2\text{R}^{11}$ ,  $-\text{NHCONR}^9\text{R}^{10}$ ,  $-\text{OR}^9$ ,  $-\text{NR}^9\text{R}^{10}$ ,  $-\text{CONR}^9\text{R}^{10}$  and  $-\text{NR}^9\text{COR}^{10}$ ; or  $B$  is  $\text{C}_{2-4}\text{alkenyl}$  or  $\text{C}_{2-4}\text{alkynyl}$ , each being optionally substituted by a group selected from  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{3-6}\text{cycloalkyl}$ , aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more halo, nitro, cyano, trifluoromethyl, trifluoromethoxy,  $-\text{CONHR}^9$ ,  $-\text{CONR}^9\text{R}^{10}$ ,  $-\text{SO}_2\text{R}^{11}$ ,  $-\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $-\text{NR}^9\text{SO}_2\text{R}^{11}$ ,  $\text{C}_{1-4}\text{alkyl}$  or
- 30  $\text{C}_{1-4}\text{alkoxy}$ ;

$R^1$  and  $R^2$  are independently hydrogen or a group selected from  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-6}$ cycloalkyl and  $C_{5-6}$ cycloalkenyl which group may be optionally substituted by halo, cyano, hydroxy or  $C_{1-4}$ alkoxy;

- $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  are independently hydrogen or a group selected from  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-6}$ cycloalkyl,  $C_{5-6}$ cycloalkenyl, aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more substituents independently selected from halo, nitro, cyano, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{3-6}$ cycloalkyl (optionally substituted by one or more  $R^{17}$ ), aryl (optionally substituted by one or more  $R^{17}$ ), heteroaryl (optionally substituted by one or more  $R^{17}$ ), heterocyclyl,  $-OR^{18}$ ,  $-SR^{19}$ ,  $-SOR^{19}$ ,  $-SO_2R^{19}$ ,  $-COR^{19}$ ,  $-CO_2R^{18}$ ,  $-CONR^{18}R^{20}$ ,  $-NR^{16}COR^{18}$ ,  $-SO_2NR^{18}R^{20}$  and  $-NR^{16}SO_2R^{19}$ ; or  $R^1$  and  $R^3$  together with the nitrogen or carbon atoms and carbon atom to which they are respectively attached form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted on carbon by  $C_{1-4}$ alkyl, fluoro or  $C_{1-4}$ alkoxy and/or on nitrogen by  $-COC_{1-3}$ alkyl,  $-SO_2C_{1-3}$ alkyl or  $C_{1-4}$ alkyl; or  $R^3$  and  $R^4$  together form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted on carbon by  $C_{1-4}$ alkyl, fluoro or  $C_{1-4}$ alkoxy and/or on nitrogen by  $-COC_{1-3}$ alkyl,  $-SO_2C_{1-3}$ alkyl or  $C_{1-4}$ alkyl; or  $R^5$  and  $R^6$  together form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted on carbon by  $C_{1-4}$ alkyl, fluoro or  $C_{1-4}$ alkoxy and/or on nitrogen by  $-COC_{1-3}$ alkyl,  $-SO_2C_{1-3}$ alkyl or  $C_{1-4}$ alkyl;
- $R^7$  is hydrogen or a group selected from  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl, heteroalkyl,  $C_{3-7}$ cycloalkyl, aryl, heteroaryl or heterocyclyl where the group is optionally substituted by halo,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $C_{3-7}$ cycloalkyl, heterocyclyl, aryl, heteroaryl or heteroalkyl; and wherein the group from which  $R^7$  may be selected is optionally substituted on the group and/or on its optional substituent by one or more substituents independently selected from halo, cyano,  $C_{1-4}$ alkyl, nitro, halo $C_{1-4}$ alkyl, heteroalkyl, aryl, heteroaryl, hydroxy $C_{1-4}$ alkyl,  $C_{3-7}$ cycloalkyl, heterocyclyl,  $C_{1-4}$ alkoxy $C_{1-4}$ alkyl, halo $C_{1-4}$ alkoxy $C_{1-4}$ alkyl,  $-COC_{1-4}$ alkyl,  $-OR^{21}$ ,  $-CO_2R^{21}$ ,  $-SR^{25}$ ,  $-SOR^{25}$ ,  $-SO_2R^{25}$ ,  $-NR^{21}COR^{22}$ ,  $-CONR^{21}R^{22}$  and  $-NHCONR^{21}R^{22}$ ;

or  $R^3$  and  $R^7$  together with the carbon atoms to which they are each attached and  $(CR^5R^6)_n$  form a saturated 5- to 7-membered ring optionally containing a heteroatom group selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted on carbon by  $C_{1-4}$ alkyl, fluoro or  $C_{1-4}$ alkoxy and/or on nitrogen by  $-COC_{1-3}$ alkyl,  $-SO_2C_{1-3}$ alkyl or  $C_{1-4}$ alkyl;

5  $R^8$  is selected from hydrogen,  $C_{1-6}$ alkyl and halo $C_{1-6}$ alkyl;

$R^9$  and  $R^{10}$  are independently hydrogen,  $C_{1-6}$ alkyl or  $C_{3-6}$ cycloalkyl;

or  $R^9$  and  $R^{10}$  together with the nitrogen to which they are attached form a heterocyclic 4 to 7-membered ring.

$R^{11}$  is  $C_{1-6}$ alkyl or  $C_{3-6}$ cycloalkyl;

10  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$  and  $R^{15}$  are independently selected from hydrogen,  $C_{1-6}$ alkyl and  $C_{3-6}$ cycloalkyl;

$R^{16}$  is hydrogen or  $C_{1-6}$ alkyl;

$R^{17}$  is selected from halo,  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl and  $C_{1-6}$ alkoxy;

$R^{18}$  is hydrogen or a group selected from  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl,  $C_{5-7}$ cycloalkenyl, saturated heterocyclyl, aryl, heteroaryl, aryl $C_{1-4}$ alkyl and heteroaryl $C_{1-4}$ alkyl which group is optionally

15 substituted by one or more halo;

$R^{19}$  and  $R^{25}$  are independently a group selected from  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl,  $C_5$ -

$\gamma$ cycloalkenyl, saturated heterocyclyl, aryl, heteroaryl, aryl $C_{1-4}$ alkyl and heteroaryl $C_{1-4}$ alkyl which group is optionally substituted by one or more halo;

$R^{20}$  is hydrogen,  $C_{1-6}$ alkyl or  $C_{3-6}$ cycloalkyl;


20 or  $R^{18}$  and  $R^{20}$  together with the nitrogen to which they are attached form a heterocyclic 4- to 7- membered ring;

$R^{21}$  and  $R^{22}$  are independently hydrogen,  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl, aryl and aryl $C_{1-4}$ alkyl;

or  $R^{21}$  and  $R^{22}$  together with the nitrogen to which they are attached form a heterocyclic 5- to 6- membered ring.

25

2. A compound of formula (1) or a pharmaceutically acceptable salt thereof:



wherein:

$Y^1$  and  $Y^2$  are independently O or S;

z is  $NR^8$ , O or S;

n is 0;

5 W is  $NR^1$ ;

m is 0 or 1;

D is hydrogen,  $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl or fluoro;

X is  $-(CR^{12}R^{13})_t-Q-(CR^{14}R^{15})_u-$  where t and u are independently 0 or 1 and Q is O, S, SO or  $SO_2$ ;

10 B is a group selected from aryl, heteroaryl and heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo, cyano,  $C_{1-4}$ alkyl (optionally substituted by  $R^9$  or  $C_{1-4}$ alkoxy or one or more halo),  $C_{2-4}$ alkenyl (optionally substituted by halo or  $R^9$ ),  $C_{2-4}$ alkynyl (optionally substituted by halo or  $R^9$ ),  $C_{3-6}$ cycloalkyl (optionally substituted by  $R^9$  or one or more halo),

15  $C_{5-6}$ cycloalkenyl (optionally substituted by halo or  $R^9$ ), aryl (optionally substituted by halo or  $C_{1-4}$ alkyl), heteroaryl (optionally substituted by halo or  $C_{1-4}$ alkyl), heterocyclyl (optionally substituted by  $C_{1-4}$ alkyl),  $-SR^{11}$ ,  $-SOR^{11}$ ,  $-SO_2R^{11}$ ,  $-SO_2NR^9R^{10}$ ,  $-NR^9SO_2R^{11}$ ,  $-NHCONR^9R^{10}$ ,  $-OR^9$ ,  $-CONR^9R^{10}$  and  $-NR^9COR^{10}$ ;

$R^1$  is hydrogen or a group selected from  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-6}$ cycloalkyl and

20  $C_{5-6}$ cycloalkenyl which group may be optionally substituted by halo, cyano, hydroxy or  $C_{1-4}$ alkoxy;

$R^3$  and  $R^4$  are independently hydrogen or a group selected from  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{3-5}$ cycloalkyl, pentenyl, aryl, heteroaryl and heterocyclyl which group is optionally substituted by one or more substituents independently selected from halo, nitro, cyano,

25 trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{3-6}$ cycloalkyl (optionally substituted by one or more  $R^{17}$ ), aryl (optionally substituted by one or more  $R^{17}$ ), heteroaryl (optionally substituted by one or more  $R^{17}$ ), heterocyclyl,  $-OR^{18}$ ,  $-SR^{19}$ ,  $-SOR^{19}$ ,  $-SO_2R^{19}$ ,  $-CONR^{18}R^{20}$  and  $-NR^{16}COR^{18}$ ;

or  $R^1$  and  $R^3$  together with the nitrogen and carbon atoms to which they are respectively

30 attached form a saturated 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and  $SO_2$  where the ring is optionally substituted on carbon

by C<sub>1-4</sub>alkyl, fluoro or C<sub>1-4</sub>alkoxy and/or on nitrogen by -COC<sub>1-3</sub>alkyl, -SO<sub>2</sub>C<sub>1-3</sub>alkyl or C<sub>1-4</sub>alkyl;

or R<sup>3</sup> and R<sup>4</sup> together form a carbocyclic or saturated heterocyclic 3- to 7-membered ring optionally containing 1 or 2 heteroatom groups selected from NH, O, S, SO and SO<sub>2</sub> where

5 the ring is optionally substituted on carbon by C<sub>1-4</sub>alkyl, fluoro or C<sub>1-4</sub>alkoxy and/or on nitrogen by -COC<sub>1-3</sub>alkyl, -SO<sub>2</sub>C<sub>1-3</sub>alkyl or C<sub>1-4</sub>alkyl;

R<sup>7</sup> is hydrogen or a group selected from C<sub>1-4</sub>alkyl, heteroalkyl, C<sub>3-5</sub>cycloalkyl, aryl, heteroaryl or heterocyclyl which group is optionally substituted by halo, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>3-5</sub>cycloalkyl, heterocyclyl, aryl, heteroaryl or heteroalkyl; and wherein the group from which R<sup>7</sup>

10 may be selected is optionally substituted on the group and/or on its optional substituent by one or more substituents independently selected from halo, cyano, C<sub>1-4</sub>alkyl, nitro, haloC<sub>1-4</sub>alkyl, heteroalkyl, aryl, heteroaryl, hydroxyC<sub>1-4</sub>alkyl, C<sub>3-5</sub>cycloalkyl, heterocyclyl, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, -COC<sub>1-4</sub>alkyl, -OR<sup>21</sup>, -CO<sub>2</sub>R<sup>21</sup>, -SR<sup>25</sup>, -SOR<sup>25</sup>, -SO<sub>2</sub>R<sup>25</sup>, -CONR<sup>21</sup>R<sup>22</sup> and -NHCONR<sup>21</sup>R<sup>22</sup>;

15 or R<sup>3</sup> and R<sup>7</sup> together with the carbon atoms to which they are each attached and (CR<sup>5</sup>R<sup>6</sup>)<sub>n</sub> form a saturated carbocyclic or heterocyclic 5- or 6-membered ring;

R<sup>8</sup> is selected from hydrogen, C<sub>1-4</sub>alkyl and haloC<sub>1-4</sub>alkyl;

R<sup>9</sup> and R<sup>10</sup> are independently hydrogen, C<sub>1-6</sub>alkyl or C<sub>3-6</sub>cycloalkyl;

or R<sup>9</sup> and R<sup>10</sup> together with the nitrogen to which they are attached form a heterocyclic 4 to 6-  
20 membered ring.

R<sup>11</sup> is C<sub>1-4</sub>alkyl or C<sub>3-5</sub>cycloalkyl;

R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> are independently selected from hydrogen, C<sub>1-4</sub>alkyl and C<sub>3-4</sub>cycloalkyl;

R<sup>16</sup> is hydrogen or C<sub>1-4</sub>alkyl;

R<sup>17</sup> is selected from halo, C<sub>1-4</sub>alkyl, C<sub>3-5</sub>cycloalkyl and C<sub>1-4</sub>alkoxy;

25 R<sup>18</sup> is hydrogen or a group selected from C<sub>1-4</sub>alkyl, C<sub>3-5</sub>cycloalkyl, C<sub>5-6</sub>cycloalkenyl, saturated heterocyclyl, aryl, heteroaryl, arylC<sub>1-4</sub>alkyl and heteroarylC<sub>1-4</sub>alkyl which group is optionally substituted by one or more halo;

R<sup>19</sup> and R<sup>25</sup> are independently a group selected from C<sub>1-4</sub>alkyl, C<sub>3-5</sub>cycloalkyl, C<sub>5-6</sub>cycloalkenyl, saturated heterocyclyl, aryl, heteroaryl, arylC<sub>1-4</sub>alkyl and heteroarylC<sub>1-4</sub>alkyl

30 which group is optionally substituted by one or more halo;

R<sup>20</sup> is hydrogen, C<sub>1-4</sub>alkyl or C<sub>3-5</sub>cycloalkyl;

or  $R^{18}$  and  $R^{20}$  together with the nitrogen to which they are attached form a heterocyclic 4- to 6- membered ring;

$R^{21}$  and  $R^{22}$  are independently hydrogen,  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl, aryl and aryl $C_{1-4}$ alkyl;

or  $R^{21}$  and  $R^{22}$  together with the nitrogen to which they are attached form a heterocyclic 5- to 6- membered ring.

3. A compound according to claim 1 wherein B is phenyl, naphthyl, pyridyl, quinolinyl, isoquinolinyl, thienopyridyl, naphthyridinyl, 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, thienopyrimidinyl, pyridoimidazolyl, benzimidazolyl, benzofuranyl, benzothienyl, indolyl, benzothiazolyl, benzotriazolyl, benzisoxazolyl, benzisothiazolyl, indazolyl, indoliziny, isobenzofuranyl, quinazolinyl, imidazopyridinyl, pyrazolopyridinyl, indolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl or isoindolinyl, where each is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo,  $C_{1-4}$ alkyl (optionally substituted by one or more halo),  $C_{2-4}$ alkynyl, heteroaryl,  $-OR^9$ , cyano,  $-NR^9R^{10}$ ,  $-CONR^9R^{10}$  and  $-NR^9COR^{10}$ ; or B is vinyl or ethynyl optionally substituted by  $C_{1-4}$ alkyl.

4. A compound according to claim 1 or 2 wherein B is a group selected from bicyclic aryl, bicyclic heteroaryl and bicyclic heterocyclyl, where each group is optionally substituted by one or more groups independently selected from nitro, trifluoromethyl, trifluoromethoxy, halo,  $C_{1-4}$ alkyl (optionally substituted by one or more halo),  $C_{2-4}$ alkynyl, heteroaryl,  $-OR^9$ , cyano,  $-NR^9R^{10}$ ,  $-CONR^9R^{10}$  and  $-NR^9COR^{10}$ ; or B is  $C_{2-4}$ alkenyl or  $C_{2-4}$ alkynyl optionally substituted by  $C_{1-4}$ alkyl,  $C_{3-6}$ cycloalkyl or heterocyclyl.

5. A compound according to claim 1 or 2 wherein B is 2-methylquinolin-4-yl.

6. A compound according to any one of the preceding claims wherein  $R^7$  is hydrogen or a group selected from  $C_{1-4}$ alkyl, aryl $C_{1-4}$ alkyl, heteroaryl $C_{1-4}$ alkyl, heterocyclyl $C_{1-4}$ alkyl, aryl, heteroaryl, heterocyclyl and  $C_{3-5}$ cycloalkyl which group is optionally substituted by cyano,  $C_{1-4}$ alkyl, halo,  $-OR^{21}$ ,  $-NR^{21}R^{22}$ ,  $-CO_2R^{21}$  and  $-NR^{21}CO_2R^{22}$ .

7. A compound according to claim 6 wherein  $R^7$  is hydrogen or  $C_{1-4}$ alkyl optionally substituted with halo, hydroxy or  $C_{1-3}$ alkoxy.
8. A pharmaceutical composition comprising a compound according to claim 1 or claim 2; and a pharmaceutically-acceptable diluent or carrier.
9. A compound according to claim 1 or 2 for use as a medicament.
10. The use of a compound according to claim 1 or 2 in the manufacture of a medicament in the treatment of a disease condition mediated  $TNF-\alpha$ .
11. The use of a compound according to claim 1 or 2 in the manufacture of a medicament in the treatment of autoimmune disease, allergic/atopic diseases, transplant rejection, graft versus host disease, cardiovascular disease, reperfusion injury and malignancy in a warm-blooded animal such as man.
12. A method of treating autoimmune disease, allergic/atopic diseases, transplant rejection, graft versus host disease, cardiovascular disease, reperfusion injury and malignancy in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound according to claim 1 or 2.
13. A process for preparing a compound according to claim 1 or 2, comprising the steps of converting a ketone or aldehyde of formula (2) into a compound of formula (1);



25

and thereafter if necessary:

- i) converting a compound of formula (1) into another compound of formula (1);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester



# INTERNATIONAL SEARCH REPORT

Int. Application No  
PC1/700 03/03937

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07D233/76 C07D401/12 A61K31/454

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	WO 02 074750 A (LUNDKVIST MICHAEL ;ASTRAZENECA AB (SE); ERIKSSON ANDERS (SE); LEPI) 26 September 2002 (2002-09-26) pages: 127-130; 159-162; 201-202 Claim 6	1, 2, 9
A	WO 00 12477 A (BRITISH BIOTECH PHARM ;MARTIN FIONNA MITCHELL (GB)) 9 March 2000 (2000-03-09) examples 5, 6 claim 1	1, 2, 9
A	WO 00 12478 A (ZENECA PHARMA SA ;TUCKER HOWARD (GB); WATERSON DAVID (GB); ZENECA) 9 March 2000 (2000-03-09) example 14 page 51, line 20 -page 52, line 2 claim 1	1, 2, 9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*G\* document member of the same patent family

Date of the actual completion of the international search

5 December 2003

Date of mailing of the international search report

19/12/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx 31 651 epo nl  
Fax (+31-70) 340-3016

Authorized officer

Fanni, S

**THIS PAGE BLANK (USPTO)**

# INTERNATIONAL SEARCH REPORT

Application No.  
PCT/GB 03/03937

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 12  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claim 12 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

---

Continuation of Box I.1

Claims Nos.: 12

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/GB 03/03937

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 02074750	A	26-09-2002	CA 2440473 A1 WO 02074767 A1 WO 02074748 A1 WO 02074750 A1 WO 02074751 A1 CA 2440475 A1 WO 02074749 A1 WO 02074752 A1	26-09-2002 26-09-2002 26-09-2002 26-09-2002 26-09-2002 26-09-2002 26-09-2002 26-09-2002
WO 0012477	A	09-03-2000	AU 5634999 A EP 1107953 A1 WO 0012477 A1 JP 2002523492 T US 2003050310 A1 US 6479502 B1	21-03-2000 20-06-2001 09-03-2000 30-07-2002 13-03-2003 12-11-2002
WO 0012478	A	09-03-2000	AU 764367 B2 AU 5524799 A BG 105369 A BR 9913255 A CA 2339761 A1 CN 1324347 T EE 200100106 A EP 1109787 A1 WO 0012478 A1 HU 0103344 A2 JP 2002523493 T NO 20011023 A NZ 509730 A PL 346344 A1 SK 2702001 A3 TR 200100605 T2 ZA 200101231 A	14-08-2003 21-03-2000 31-12-2001 22-05-2001 09-03-2000 28-11-2001 17-06-2002 27-06-2001 09-03-2000 28-02-2002 30-07-2002 25-04-2001 30-05-2003 11-02-2002 06-08-2001 21-08-2001 13-05-2002

**THIS PAGE BLANK (USPTO)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**



**THIS PAGE BLANK (USPTO)**